

THE SCHRÖDINGER PROPAGATOR FOR SCATTERING METRICS

ANDREW HASSELL AND JARED WUNSCH

ABSTRACT. Let g be a scattering metric on a compact manifold X with boundary, i.e., a smooth metric giving the interior X° the structure of a complete Riemannian manifold with asymptotically conic ends. An example is any compactly supported perturbation of the standard metric on \mathbb{R}^n . Consider the operator $H = \frac{1}{2}\Delta + V$, where Δ is the positive Laplacian with respect to g and V is a smooth real-valued function on X vanishing to second order at ∂X . Assuming that g is non-trapping, we construct a global parametrix $\mathcal{U}(z, w, t)$ for the kernel of the Schrödinger propagator $U(t) = e^{-itH}$, where $z, w \in X^\circ$ and $t \neq 0$. The parametrix is such that the difference between \mathcal{U} and U is smooth and rapidly decreasing both as $t \rightarrow 0$ and as $z \rightarrow \partial X$, uniformly for w on compact subsets of X° . Let $r = x^{-1}$, where x is a boundary defining function for X , be an asymptotic radial variable, and let $W(t)$ be the kernel $e^{-ir^2/2t}U(t)$. Using the parametrix, we show that $W(t)$ belongs to a class of ‘Legendre distributions’ on $X \times X^\circ \times \mathbb{R}_{\geq 0}$ previously considered by Hassell-Vasy. When the metric is trapping, then the parametrix construction goes through microlocally in the non-trapping part of the phase space.

We apply this result to determine the singularities of $U(t)f$, for any tempered distribution f and for any fixed $t \neq 0$, in terms of the oscillation of f near ∂X . If the metric is nontrapping then we *precisely* determine the wavefront set of $U(t)f$, and hence also precisely determine its singular support. More generally, we are able to determine the wavefront set of $U(t)f$ for $t > 0$, resp. $t < 0$ on the non-backward-trapped, resp. non-forward-trapped subset of the phase space. This generalizes results of Craig-Kappeler-Strauss and Wunsch.

1. INTRODUCTION

Let (X, g) be a scattering manifold of dimension n . Thus, X is a compact n -dimensional manifold with boundary, and g is a metric in the interior of X taking the form

$$g = \frac{dx^2}{x^4} + \frac{h}{x^2}$$

near the boundary. Here x is a boundary defining function, y are local coordinates on the boundary $Y = \partial X$ extended to X and h is a 2-cotensor that restricts to a metric (i.e., is positive definite) on Y . We shall assume that x is a globally defined, smooth function on X that vanishes only at ∂X , and denote $r = x^{-1}$, which is analogous to the radial variable on Euclidean space. The metric takes a more familiar form when written in terms of r :

$$g = dr^2 + r^2h(r^{-1}, y, dr, dy),$$

where h is smooth in the first variable and y (smoothness in $x = 1/r \in [0, \epsilon]$ is of course a much stronger condition than smoothness in $r \in (\epsilon^{-1}, \infty)$). In fact, one can choose coordinates locally so that g takes the form

$$(1.1) \quad g = dr^2 + r^2h(r^{-1}, y, dy) = \frac{dx^2}{x^4} + \frac{h(x, y, dy)}{x^2};$$

This research was supported in part by a Fellowship and a Linkage grant from the Australian Research Council (A. H.) and by NSF grant DMS-0100501 (J. W.).

here h is a metric in the y variables depending parametrically (and smoothly) on x (see [11] for a proof). Thus, g is *asymptotically conic*; it approaches the conic metric $dr^2 + r^2h_0$, where $h_0 = h(0, y, dy)$, as $r \rightarrow \infty$. The boundary ∂X is then at geometric infinity, with each point of ∂X representing an asymptotic direction of geodesics. Euclidean space, with its standard metric or any short range perturbation of it, is an example. We shall assume henceforth that coordinates have been chosen so that the representation (1.1) for the metric holds. Let Δ denote the positive Laplacian with respect to g , and let $H = \frac{1}{2}\Delta + V$, where $V \in x^2\mathcal{C}^\infty(X; \mathbb{R})$. Thus V is a short-range potential¹. We consider the time-dependent Schrödinger equation

$$(1.2) \quad (D_t + H)u(z, t) = 0, \quad u(z, 0) = f(z) \in C^{-\infty}(X),$$

where $z \in X$, $t \in \mathbb{R}$, $D_t = -i\frac{\partial}{\partial t}$ and f is a given distribution on X . Let $U(t) = e^{-itH}$ be the propagator for H . We wish to construct a parametrix $\mathcal{U}(t)$ for $U(t)$ which captures all the singularities of U ; in particular, we want $\mathcal{U}(t) - U(t)$ to have a kernel which vanishes rapidly both as $t \rightarrow 0$ and as we approach ∂X . However, to simplify the construction we shall let only one of the variables $(z, w) \in X^2$, say the left variable z , approach infinity, while we shall only require uniformity over compact subsets in the w variable.

To state our main theorem, we choose a function $\phi \in \mathcal{C}^\infty(X)$ which is zero in a neighbourhood of the boundary of X , we let $r = x^{-1}$ as above, and we define the kernel $W(t) = e^{-ir^2/2t}U(t)$. We remark that we regard $U(t)$ and $W(t)$ as acting on half-densities, so the kernels contain a Riemannian half-density factor $|dg_z dg_w dt|^{1/2}$ in each variable. Then our main result is

Theorem 1.1. *Assume that the metric g is non-trapping. Then the kernel $\phi(w)W(t)$ is a fibred-scattering Legendrian distribution on $X \times X^\circ \times \mathbb{R}_+$ of order $(\frac{3}{4}, \frac{1}{4})$ in the sense of [8], associated to the Legendre submanifold ${}^\phi L$ defined in Lemma 3.5.*

Remark. In the trapping case, one can still construct a parametrix microlocally in the non-trapping region; see Section 5.

In more prosaic language, Theorem 1.1 says that the propagator is given by oscillatory integrals of certain rigidly prescribed forms. Near $t = 0$, and with both variables w, z away from the boundary of X , these take the form of a Legendrian distribution (see Melrose-Zworski [15])

$$t^{-n/2-k/2}(2\pi)^{-k} \int_K e^{i\Phi(z, w, v)/t} a(z, w, v, t) dv,$$

where the integral is over a compact set $K \subset \mathbb{R}^k$. Here, Φ and a are smooth in all their variables. Near the diagonal, Φ is given by $\text{dist}(w, z)^2/2$, as in the flat Euclidean case, no v variables are required, and there is no integral. Associated to Φ is a Legendrian submanifold of $T^*X \times T^*X \times \mathbb{R}$, here given explicitly by

$$L = \{(w, \xi, z, \zeta, \tau) \mid (z, \zeta) = \exp_g(w, \xi), \tau = \frac{|\xi|^2}{2}\}$$

(this is a somewhat non-invariant description, for precise details see §3). The function Φ becomes non-smooth outside the injectivity radius; geometrically this corresponds to the Legendrian L becoming non-projectable (i.e. the projection from the Legendrian to the base $X \times X$ is no longer a diffeomorphism). The Legendrian, however, remains perfectly smooth; it is no longer parametrized by a function just of w and z , but one needs extra variables v

¹Gravitational long-range potentials and metrics in the sense of [7] can also be treated; see [9].

(precisely, one needs at least k extra variables locally where k is the dimension of the null space of the differential of the projection from the Legendrian to the base).

When z approaches the boundary of X , we use local coordinates (x, y) as described above, and sometimes use $r = x^{-1}$, the asymptotic ‘radial’ variable. In this region the propagator takes a more complicated form. We show that $W(t)$ is a finite sum of oscillatory integrals of the form

$$(1.3) \quad t^{-n/2-k/2}(2\pi)^{-k} \int_K e^{i\psi(x,y,w,v)/xt} a(x, y, w, v, t) dv,$$

where again ψ and a are smooth in all their variables, and one integrates over a compact set $K \subset \mathbb{R}^k$. Notice that ψ depends on all variables apart from t .

Next, we address the question of determining the wavefront set of $u(\cdot, t)$ at a fixed nonzero (say, positive) time t in terms of the wavefront set of the initial data f . We are particularly interested in the case of ‘interior singularities’, lying above a point in the interior of X . It is well known that equation (1.2) has infinite speed of propagation, so to answer this question we must look for singularities of f at the boundary of X . Dually, since $U(t) = U(-t)^*$, we can consider interior singularities of f and determine the boundary singularities of $u(t)$ that they produce. It is known from [22] that if there is an interior singularity (w, η) for f , then for all positive times t there are quadratic oscillations of frequency $1/2t$ in $u(\cdot, t)$ at infinity in the asymptotic direction of the geodesic emanating from (w, η) . Roughly speaking, the quadratic oscillations look like $e^{ir^2/2t}$ in a conic neighbourhood of the asymptotic direction y ; more precisely, this result is phrased in terms of the ‘quadratic scattering wavefront set’ (see [22]).

The limitation of the result of [22] is that different singularities along a single geodesic, or along a different geodesic with the same asymptotic direction, produce identical quadratic oscillation, hence consideration of the quadratic wavefront set alone will not result in precise propagation results. To analyse the finer structure of $u(\cdot, t)$, we divide $u(\cdot, t)$ by the explicit quadratic oscillatory factor $e^{ir^2/2t}$, and find that the resulting function has oscillations which are *linear* in r that contain the desired information on the location of the interior singularity. The presence of linear oscillations of a function f is measured by the *scattering wavefront set* $\text{WF}_{\text{sc}}(f)$ (see [16]), a closed subset of ${}^{\text{sc}}T_{\partial X}^*X$. (This bundle, defined in §2, is the restriction to ∂X of an n -dimensional bundle over X which is a compressed and scaled version of the cotangent bundle T^*X .) We show that the asymptotics of geodesic flow on X determine two contact transformations S_f and S_b , which we call the forward and backward sojourn relations, from the sets \mathbf{F} , resp. $\mathbf{B} \subset S^*X^\circ$ consisting of points in the cosphere bundle which are not forward, resp. backward trapped under geodesic flow, to ${}^{\text{sc}}T_{\partial X}^*X$. They are related by $\mathbf{F} = -\mathbf{B}$ and $S_f(\zeta) = -S_b(-\zeta)$. The definition of S_f in local coordinates is as follows: Let $\gamma(s)$ be the arclength-parametrized geodesic emanating from $(w, \eta) \in S^*X^\circ$, let $y = (y_1, \dots, y_{n-1})$ be local coordinates on ∂X and let $(y, \nu, \mu = (\mu_1, \dots, \mu_{n-1}))$ be the induced coordinates on ${}^{\text{sc}}T_{\partial X}^*X$ as in (2.1). Then $S_f(w, \eta) = (y_0, \nu, \mu)$ iff

$$y_0 = \lim_{s \rightarrow +\infty} \gamma(s) \in \partial X$$

is the asymptotic direction of the geodesic,

$$(1.4) \quad \nu = \lim_{s \rightarrow +\infty} (s - r(\gamma(s))), \quad r = \frac{1}{x}$$

is the ‘sojourn time’ of the geodesic, and

$$(1.5) \quad \mu_i = \sum_j h_{ij} \lim_{s \rightarrow +\infty} (y_0 - y(\gamma(s)))_j / x$$

measures its angle of approach to ∂X . (That the point $y_0 \in \partial X$ should be thought of as the asymptotic *direction* of the geodesic can be seen easily in the special case when X is the radial compactification of \mathbb{R}^n .) Note that the sojourn time, thought of as depending on a geodesic and a point along it, is closely related to the classical Busemann function of differential geometry; see for example [18], chapter 1, section 2. We use the term ‘sojourn relation’ for S_f since the coordinate ν is analogous to the sojourn time considered by Guillemin [5].

Theorem 1.2. *Let f be a tempered distribution on X and set $u(\cdot, t) = e^{-itH} f$. Let $\zeta = (w, \eta) \in \mathbf{B} \subset S^* X^\circ$ be a non-backward-trapped point of $S^* X^\circ$. For any fixed $t > 0$,*

$$(1.6) \quad \zeta \in \text{WF}(u(\cdot, t)) \text{ if and only if } \frac{1}{t} S_b(\zeta) \in \text{WF}_{\text{sc}}(e^{ir^2/2t} f).$$

Here the factor t^{-1} acts by scaling the fibre variables. Similarly, if $\zeta \in \mathbf{F}$, then for $t < 0$,

$$(1.7) \quad \zeta \in \text{WF}(u(\cdot, t)) \text{ if and only if } \frac{1}{|t|} S_f(\zeta) \in \text{WF}_{\text{sc}}(e^{ir^2/2t} f).$$

Remark. The condition on the right hand side of (1.6) or (1.7) is not manifestly coordinate invariant. The sojourn relation changes under a change of coordinates, but so does the scattering wavefront set of $e^{ir^2/2t} f$, in such a way that the condition as a whole is coordinate invariant. A manifestly invariant description may be given in terms of the affine bundle of Lemma 3.3.

Example. Consider the free Hamiltonian $H = \frac{1}{2} \Delta$ on \mathbb{R}^n . The propagator in this case is

$$U(t) = (2\pi t)^{-n/2} e^{i|z-w|^2/2t}, \quad \text{whence} \quad W(t) = (2\pi t)^{-n/2} e^{-iz \cdot w/t} e^{i|w|^2/2t}.$$

It is not hard to check that $W(t)$ satisfies the conclusion of Theorem 1.1 in this case. Consider initial data

$$f = (2\pi T)^{-n/2} e^{-i|z-w|^2/2T}, \quad T > 0.$$

Then the solution at time $t = T$ is a delta function centred at w , hence its wavefront set is

$$\{(w, \hat{z}) \mid \hat{z} \in S^{n-1}\}.$$

In the free case, the backward sojourn relation is given by

$$S_b(w, \hat{z}) = (y = -\hat{z}, \nu = -w \cdot \hat{z}, \mu = w - (w \cdot \hat{z})\hat{z}),$$

and we easily check that (1.6) holds in this case.

Most previous work on Schrödinger parametrices has focused on the case of flat space with a potential perturbation, where the geometric situation is substantially simpler. Very detailed parametrix constructions in this setting have been made by Fujiwara [4], Zelditch [24], Trèves [21] and Yajima [23]. In the case of curved space, very little was known. Kapitanski-Safarov [13] have shown that on \mathbb{R}^n with a compactly-supported, nontrapping potential perturbation, the fundamental solution is smooth for $t > 0$, and have exhibited a parametrix modulo $\mathcal{C}^\infty(\mathbb{R}^n)$ [12]. Such a parametrix, however, is not sufficiently specified at infinity to yield results about smoothness of the solution of the general Cauchy problem at $t > 0$.

Regularity results for the Schrödinger propagator for non-flat metrics in the form of Strichartz estimates have been obtained recently by Staffilani-Tataru [19] and Burq-Gérard-Tzvetkov [1]. Staffilani and Tataru [19] proved Strichartz estimates for $e^{-itH}f$ where H is the Laplacian of a C^2 , compactly supported, nontrapping perturbation of the standard metric on \mathbb{R}^n , using the FBI transform and Littlewood-Paley decompositions to handle the rough metric. Burq, Gérard and Tzvetkov obtained Strichartz estimates, with a loss of derivatives compared to the flat Euclidean case, for compact manifolds or perturbations of the Laplacian on \mathbb{R}^n , without any nontrapping assumption.

Various authors have considered the question of determining the singularities of $u = e^{-itH}f$ in terms of f . The first results about microlocal smoothness of $u(t)$ for $t > 0$ and general initial data were those of Craig-Kappeler-Strauss [3], who showed that, on an asymptotically Euclidean space, decay of the initial data in a microlocal incoming cone yields microlocal smoothness along the whole pencil of geodesics emanating from that cone for all $t > 0$. This result was refined in [22], where the second author showed that absence of the quadratic-scattering wavefront set (see Section 2.2) allowed one to conclude absence at varying *times* and along varying pencils of geodesics. This approach, while it specified in terms of the Cauchy data *when* and *in what direction* singularities might appear in X° , failed to say anything about *where* they might land. These propagation results have been extended to the analytic category by Robbiano-Zuily [17].

Our Theorem 1.2 gives a complete solution to the propagation problem in the case when the metric g is nontrapping, and in general, a complete characterization of the singularities of $u(t)$ in the non-backward-trapped set \mathbf{F} for $t > 0$, and in the non-forward-trapped set \mathbf{B} for $t < 0$. Our results imply those of [22], and hence those of [3]², since the hypothesis on f required in [22] for microlocal smoothness of $u(t)$ along all geodesics emanating from $y \in \partial X$ implies that in fact $(y, \nu, \mu) \notin \text{WF}_{\text{sc}} e^{i\nu^2/2t} f$ for all (ν, μ) . Hence Theorem 1.2 yields the main boundary to interior propagation result of [22] as a special case.

We thank Richard Melrose, András Vasy and Steve Zelditch for useful conversations, and the Erwin Schrödinger Institute, the Mathematics Department at SUNY Stonybrook and the Mathematical Sciences Institute at the Australian National University for their hospitality.

2. CONTACT STRUCTURES AND LEGENDRIAN DISTRIBUTIONS

We recall the definition of various structures associated to manifolds with boundary and corners needed in this paper. For further details, see Melrose [16], Melrose-Zworski [15], Hassell-Vasy [7, 8].

2.1. Scattering structure and Legendrian distributions. Let X be a d -dimensional manifold with boundary, and let x be a boundary defining function for X . We identify a collar neighborhood of ∂X with $\partial X \times [0, \epsilon]$ so that specifying y_1, \dots, y_{d-1} local coordinates in ∂X gives local coordinates (x, y) for X . The Lie algebra of scattering vector fields $\mathcal{V}_{\text{sc}}(X)$ consists of vector fields of the form $ax^2\partial_x + \sum xb_i\partial_{y_i}$ with $a, b_i \in C^\infty(X)$. Such vector fields can be described as the set of C^∞ sections of a vector bundle ${}^{\text{sc}}TX$. The dual of this bundle we denote ${}^{\text{sc}}T^*X$; sections of it are locally spanned over $C^\infty(X)$ by dx/x^2 and dy_i/x . Hence any point $q \in {}^{\text{sc}}T^*_{x,y}X$ has a unique expression

$$(2.1) \quad q = \nu d\left(\frac{1}{x}\right) + \sum \mu_i \frac{dy_i}{x}$$

²although we require more decay of our potential than is assumed in [3].

which yields local coordinates (x, y, ν, μ) for ${}^{\text{sc}}T^*X$, ν and μ being linear on each fibre. We say that a half-density α on X is a scattering half-density if it is a smooth and non-vanishing section of the bundle $(\wedge^n({}^{\text{sc}}T^*X))^{1/2}$; such sections have the form $a|dx dy_1 \dots dy_{d-1}/x^{d+1}|^{1/2}$, where $a \in \mathcal{C}^\infty(X)$ is nonvanishing. The restriction of ${}^{\text{sc}}T^*X$ to the boundary of X is denoted ${}^{\text{sc}}T_{\partial X}^*X$.

We define $\dot{\mathcal{C}}^\infty(X) = \bigcap_{l \geq 0} x^l \mathcal{C}^\infty(X)$, with its natural Fréchet topology, and denote by $\mathcal{C}^{-\infty}(X)$ its topological dual. We sometimes refer to these as the space of Schwartz functions and the space of tempered distributions on X , by analogy with Euclidean space.

Now we recall some facts about the scattering calculus $({}^{\text{sc}}\Psi^{m,l}(X; {}^{\text{sc}}\Omega^{1/2}(X))$ acting on half-densities, which is indexed by two orders (m, l) , the interior order m (which for a differential operator is the order of the highest derivative that occurs) and the boundary order l . The half-density factor will be understood from now on, and dropped from notation. The space $\Psi^{m,l}(X)$ is the same as $x^l \Psi^{m,0}(X)$, and $\Psi^{m,0}(X)$ is a ‘microlocalization’ of the scattering differential operators of order m on X ; it contains in particular all m th order differential operators generated over $\mathcal{C}^\infty(X)$ by $\mathcal{V}_{\text{sc}}(X)$. Operators $P \in {}^{\text{sc}}\Psi^{m,l}(X)$ are determined up to ${}^{\text{sc}}\Psi^{m,l+1}(X)$ by the boundary symbol p , which is a smooth function on the boundary of the scattering cotangent bundle, ${}^{\text{sc}}T_{\partial X}^*X$, and up to ${}^{\text{sc}}\Psi^{m,l+2}(X)$ by the boundary symbol p together with the boundary subprincipal symbol, which is again a smooth function p_{sub} on ${}^{\text{sc}}T_{\partial X}^*X$. The operator P is said to be elliptic at $q \in {}^{\text{sc}}T_{\partial X}^*X$ if $p(q) \neq 0$. The scattering wavefront set $\text{WF}_{\text{sc}}(u)$ of a distributional half-density u is defined by the condition that $q \in {}^{\text{sc}}T_{\partial X}^*X$ is *not* in $\text{WF}_{\text{sc}}(u)$ iff there is an $A \in {}^{\text{sc}}\Psi^{0,0}(X)$ such that A is elliptic at q and $Au \in \dot{\mathcal{C}}^\infty(X)$, the space of Schwartz functions on X° . The scattering wavefront set of u is always a closed subset of ${}^{\text{sc}}T_{\partial X}^*X$. We also have a scale of Sobolev spaces $H_{\text{sc}}^m(X)$, defined by $u \in H_{\text{sc}}^m(X)$ iff $V_1 \cdots V_m u \in L^2(X)$ for all $V_1, \dots, V_m \in \mathcal{V}_{\text{sc}}(X)$, or equivalently, if $Pu \in L^2(X)$ for all $P \in \Psi^{m,0}(X)$. (Here $L^2(X)$ is defined with respect to the Riemannian measure dg .)

For the purposes of this paper, we will often take the manifold with boundary to be $X^\circ \times X^\circ \times \mathbb{R}_{\geq 0}$ (this space is not compact, but that is irrelevant here). In that case the boundary defining function is t and local coordinates on the boundary will be denoted (z, w) , where $z \in \mathbb{R}^n$ is a local coordinate for the first factor and $w \in \mathbb{R}^n$ is a local coordinate for the second. In this case we use coordinates $(t, z, w, \tau, \zeta, \eta)$ where we write points $q' \in {}^{\text{sc}}T^*(X^\circ \times X^\circ \times \mathbb{R}_{\geq 0})$

$$(2.2) \quad q' = \tau d\left(\frac{1}{t}\right) + \sum_{i=1}^n \zeta_i \frac{dz_i}{t} + \sum_{j=1}^n \eta_j \frac{dw_j}{t}.$$

Returning to the general situation, there is a contact structure defined at the boundary ${}^{\text{sc}}T_{\partial X}^*X$ of ${}^{\text{sc}}T^*X$. It is defined by the contact one-form

$$(2.3) \quad \chi = \omega(x^2 \partial_x, \cdot) \lrcorner \{x = 0\},$$

where ω is the symplectic form on T^*X (which is canonically isomorphic to ${}^{\text{sc}}T^*X$ over X°). In local coordinates $\chi = \sum \mu_i dy_i - d\nu$, so χ is clearly nondegenerate. A change of boundary defining function $x \rightarrow ax$ changes χ according to $\chi \rightarrow a\chi$, so the contact structure defined by χ is completely natural. A Legendrian submanifold of ${}^{\text{sc}}T_{\partial X}^*X$ is defined, as usual, to be a smooth submanifold of maximal dimension, namely $\dim X - 1$, such that the contact form χ vanishes on it. Any Legendrian submanifold L has a local nondegenerate parametrization in a neighbourhood of any $q \in L$. This, by definition, is a function $\psi(y, v)$, with $v \in \mathbb{R}^k$ for

some $k \geq 0$, such that

$$(2.4) \quad \begin{aligned} &\text{the differentials } d\left(\frac{\partial\psi}{\partial v_i}\right), \quad i = 1, \dots, k \text{ are linearly independent whenever } d_v\psi = 0, \\ &\text{and } L = \{(y, d(\frac{\psi}{x})) \mid d_v\psi = 0\} \text{ locally near } q. \end{aligned}$$

The simplest situation is when L is projectable in the sense that the projection $(y, \nu, \mu) \rightarrow y$ from ${}^{\text{sc}}T_{\partial X}^*X$ to ∂X restricts to a diffeomorphism from L to ∂X . Then y is a coordinate on L , so ν is given by a function $\psi(y)$ on L . In this case, no extra variables v are required, and L is given by $\{(y, d(\psi/x))\}$ locally. If the kernel of the differential of the projection $(y, \nu, \mu) \rightarrow y$ restricted to L has dimension k then at least k extra variables are required to locally parametrize L near q .

A (half-density) Legendrian distribution of order m associated to L is a half-density $u\alpha$, where α is a scattering half-density and u is a finite sum of terms $\sum_i u_i + u_0$, where $u_0 \in \dot{C}^\infty(X)$ and u_i is given by an certain type of oscillatory integral associated to a local parametrization of L :

$$(2.5) \quad u_i = (2\pi)^{-k} x^{m-k/2+d/4} \int_K e^{i\psi(y,v)/x} a(y, v, x) dv.$$

Here a is a smooth function of x, y, v with compact support and ψ is a nondegenerate parametrization of L on the support of a . The set of half-density Legendrian distributions of order m associated with L is denoted $I^m(X, L; {}^{\text{sc}}\Omega^{1/2}(X))$, or just $I^m(L)$ when the space X is understood (in this paper, Legendrian distributions will always be half-densities).

Legendrian distributions have a well-defined symbol map σ^m taking values in smooth sections of a line bundle $S^{[m]}(L)$ over L . This bundle is given by $S^{[m]}(L) = {}^{\text{sc}}\Omega^{1/2}(L) \otimes |N^*(\partial X)|^{m-d/4} \otimes M \otimes E$, where $\Omega^{1/2}(L)$ denotes the half-density bundle over L , N^*X is the conormal bundle, M is the Maslov bundle and E is the bundle described in [7]. To define the symbol, we choose $d-1$ functions λ_j in (y, v) -space which together with $d_{v_i}\psi$ give local coordinates in (y, v) -space. Then $\{\lambda_j\}$ are local coordinates on L via the identification (2.4). The symbol is given, using the identification (2.4) and up to Maslov factors, by

$$(2.6) \quad \sigma^m(u) = a(0, y, v) \left| \det \frac{\partial(\lambda, \partial_v\psi)}{\partial(y, v)} \right|^{-1/2} |d\lambda|^{1/2} \upharpoonright \{\partial\psi/\partial v = 0\}.$$

The symbol of $u \in I^m(L)$ determines u modulo $I^{m+1}(L)$. Consideration of how the symbol changes under changes of parametrization ψ , changes of coordinates (x, y) , and changes of coordinates λ , show that the symbol lives in the bundle $S^{[m]}(L)$ above [7]. If L is locally projectable, then the situation simplifies. We may take coordinates λ on L to be y , and the v variables are absent, the determinant factor above is 1 and the symbol becomes

$$a(0, y) |dy|^{1/2}.$$

The important bundle in the factorization of $S^{[m]}(L)$ here is $|N^*(\partial X)|^{m-d/4}$; in particular this tells us that if we change boundary defining function from x to $xa^{-1}(y)$ then a symbol of order m changes by a factor $a^{m-d/4}$.

We now recall the symbol calculus for a scattering pseudodifferential operator $P \in {}^{\text{sc}}\Psi^{*,0}(X)$ acting on a Legendrian distribution $u \in I^m(L)$. In fact, we only need to consider the case when the symbol p of P vanishes identically on L , and in view of our application in §4, we use coordinates (2.2) appropriate to the manifold $M = X^\circ \times X^\circ \times \mathbb{R}_{\geq 0}$. Then the Hamilton vector field of the function p (extended into the interior of ${}^{\text{sc}}T^*M$ arbitrarily),

H_p , vanishes to first order at the boundary of ${}^{\text{sc}}T^*M$, so we define the rescaled Hamilton vector field ${}^{\text{sc}}H_p$ to be $x^{-1}H_p$ restricted to ${}^{\text{sc}}T^*_\partial M M$. Then $Pu \in I^{m+1}(L)$ with symbol

$$(2.7) \quad \sigma^{m+1}(Pu) = \left(-i\mathcal{L}^{\text{sc}}_{H_p} - i\left(\frac{1}{2} + m - \frac{d}{4}\right)\frac{\partial p}{\partial \tau} + p_{\text{sub}} \right) \sigma^m(u) \otimes |dt|.$$

Here $d = 2n + 1$ is the dimension of M , and \mathcal{L}_{H_p} denotes the Lie derivative acting on half-densities. The operation of tensoring with $|dt|$ is simply the natural isomorphism between $S^{[m]}(L)$ and $S^{[m+1]}(L)$.

2.2. Quadratic scattering structure. We only touch on this very briefly since we only make fleeting use of it in this paper. The quadratic-scattering Lie algebra is defined by $\mathcal{V}_{\text{qsc}} = x\mathcal{V}_{\text{sc}}(X)$. It is the space of smooth sections of a vector bundle ${}^{\text{qsc}}TX$ which is locally spanned (near the boundary) by $x^3\partial_x$ and $x^2\partial_{y_i}$. The dual bundle is denoted ${}^{\text{qsc}}T^*X$; sections of this bundle are locally spanned over $\mathcal{C}^\infty(X)$ by dx/x^3 and dy_i/x^2 . The vector fields in $\mathcal{V}_{\text{qsc}}(X)$ lie in a calculus of quadratic scattering pseudodifferential operators much like the scattering calculus, and with a boundary symbol and associated wavefront set lying in ${}^{\text{qsc}}T^*_{\partial X}X$. It is in terms of this wavefront set that the propagation results of [22] are couched, but it will not directly concern us here.

2.3. Manifolds with corners with fibred boundaries. Above, we have looked at $X^\circ \times X^\circ \times \mathbb{R}_{\geq 0}$. However, in our analysis of the Schrödinger propagator $U(z, w, t)$, we wish to let the first variable z approach the boundary of X . Hence we need to study the manifold

$$(2.8) \quad X_t^2 = X \times X^\circ \times \mathbb{R}_{\geq 0}.$$

We denote the two boundary hypersurfaces of X_t^2 at $t = 0$ and at $x = 0$ by mf and sf, the ‘main face’ and the ‘side face’, respectively, and we denote the corner $\text{mf} \cap \text{sf}$ by K . Thus $\text{mf} = X \times X^\circ$, $\text{sf} = \partial X \times X^\circ \times \mathbb{R}_{\geq 0}$ and $K = \partial X \times X^\circ$. We shall see that the geometry of this manifold arising from the structure of the Schrödinger operator $D_t + H$ is that of a manifold with corners with fibred boundaries; see [7] and [8] for a discussion of the general situation. Here, we restrict ourselves to describing the particular situation of X_t^2 , which is considerably simpler.

We first describe the relevant fibration. Define $\phi : \text{sf} \rightarrow \partial X \times X^\circ$ by projection off the last factor:

$$(2.9) \quad \phi(y, w, t) = (y, w) \in \partial X \times X^\circ.$$

We then define a Lie algebra ${}^\phi\mathcal{V}$ of smooth vector fields on X_t^2 as follows: if V is a smooth vector field on X_t^2 , then

$$(2.10) \quad V \in {}^\phi\mathcal{V} \text{ if and only if } V(xt) = O(x^2t^2), \text{ } V \text{ is tangent to} \\ \text{the fibration } \phi \text{ at } x = 0 \text{ and } V \text{ vanishes at } t = 0.$$

This Lie algebra is independent of the choice of boundary defining function x . Corresponding to this Lie algebra is a vector bundle ${}^\phi TX_t^2$ such that ${}^\phi\mathcal{V}$ is canonically identified with the space of smooth sections of ${}^\phi TX_t^2$. In local coordinates, this is spanned (near the boundary $x = 0$) by

$$(2.11) \quad xt\partial_{y_i}, \quad xt\partial_{w_i}, \quad t(t\partial_t - x\partial_x), \quad tx^2\partial_x.$$

Note that $t^2\partial_t$ and $tx\partial_x$ are not separately in ${}^\phi\mathcal{V}$ (since they do not satisfy (2.10)), but their difference is. The dual bundle ${}^\phi T^*X_t^2$ is spanned over $\mathcal{C}^\infty(X_t^2)$ by the one-forms

$$d\left(\frac{1}{xt}\right) = -\frac{xdt + tdx}{x^2t^2}, \quad d\left(\frac{1}{t}\right), \quad \frac{dy_i}{xt}, \quad \frac{dw_i}{xt}.$$

which pair non-degenerately with (2.11). Thus every point in $\phi T^* X_t^2$ near $\{x = 0\}$ may be expressed

$$\sigma d\left(\frac{1}{xt}\right) + \tau d\left(\frac{1}{t}\right) + \bar{\mu} \frac{dy_i}{xt} + \bar{\xi} \frac{dw_i}{xt}$$

($\bar{\mu}$ and $\bar{\xi}$ are used since we are reserving the symbols μ and ξ for later use), which yields local coordinates $(x, y, w, t, \sigma, \bar{\mu}, \bar{\xi}, \tau)$ on $\phi T^* X_t^2$. A fibred-scattering half-density is defined to be a section of $\bigwedge^{2n+1}(\phi T^* X_t^2)$, $n = \dim X$, which is smooth and nonvanishing. Near the boundary, such a half-density has the form

$$a \left| \frac{dx dy dw dt}{t^{2n+2} x^{2n+1}} \right|^{1/2}, \text{ where } a \in \mathcal{C}^\infty(X_t^2) \text{ is everywhere nonzero.}$$

On $\phi T_{\text{mf}}^* X_t^2$, (that is, at $t = 0$), $\omega(xt^2 \partial_t, \cdot)$ is a contact form (i.e. nondegenerate) for $x > 0$, but it degenerates at $x = 0$. This is evident from its local coordinate expression

$$\omega(xt^2 \partial_t, \cdot) \upharpoonright \{t = 0\} = d\sigma + x d\tau + \bar{\mu} dy + \bar{\xi} dw.$$

As described in [8], although this is degenerate at $\phi T_K^* X_t^2$, that is, at $\phi T_{\text{mf}}^* X_t^2 \cap \{x = 0\}$, there is a natural fibration β from $\phi T_K^* X_t^2$ to a bundle γ over K , with a nondegenerate contact form on γ such that $\omega(xt^2 \partial_t, \cdot)$ restricted to $\{t = 0, x = 0\}$ is the lift of the contact form from γ to $\phi T_K^* X_t^2$. Here the bundle γ is naturally isomorphic to ${}^{\text{sc}}T_{\partial X \times X^\circ}^*(X \times X^\circ)$, the contact form is the natural contact form (2.3) on ${}^{\text{sc}}T_{\partial X \times X^\circ}^*(X \times X^\circ)$ and the fibration β is given by

$$(2.12) \quad \beta(0, y, w, 0, \sigma, \bar{\mu}, \bar{\xi}, \tau) = (y, w, \sigma, \bar{\mu}, \bar{\xi})$$

in the coordinates above.

A fibred-scattering Legendrian submanifold J of $\phi T^* X_t^2$, as defined in [8], is a submanifold of $\phi T_{\text{mf}}^* X_t^2$ which is Legendrian with respect to the contact form $\omega(xt^2 \partial_t, \cdot)$ for $x > 0$, which meets the boundary of $\phi T_{\text{mf}}^* X_t^2$ at $x = 0$ transversally, and such that the fibration β , restricted to ∂J , is a diffeomorphism from ∂J to a Legendrian submanifold G of ${}^{\text{sc}}T_{\partial X \times X^\circ}^*(X \times X^\circ)$ (with respect to the natural contact structure on ${}^{\text{sc}}T_{\partial X \times X^\circ}^*(X \times X^\circ)$).

A local nondegenerate parametrization of J near $q \in \partial J$, is a function $\psi(x, y, w, v)$, with $v \in \mathbb{R}^k$ for some $k \geq 0$, such that, locally near q ,

$$(2.13) \quad \begin{aligned} &\text{the differentials } d\left(\frac{\partial \psi}{\partial v_i}\right), \quad 1 \leq i \leq k \text{ are linearly independent} \\ &\text{and } J = \left\{ \left(y, d\left(\frac{\psi}{xt}\right) \right) \mid d_v \psi = 0 \right\}. \end{aligned}$$

It then follows that the function $\psi(0, y, w, v)$ parametrizes the Legendrian G .

A Legendrian distributional half-density of order (m, r) associated to J is a half-density $u\tilde{\alpha}$, where $\tilde{\alpha}$ is a fibred-scattering half-density and u is a finite sum of terms $\sum_j u_j + u_0 + u_f$, where u_0 is a Legendrian distribution of order m associated to the interior of J (where the structure is locally of scattering type, so this has already been defined), u_f is a fibred Legendrian distribution of order r associated to G , which is supported away from mf (defined below), and u_j is given by an certain type of oscillatory integral associated to a local parametrization of J . Namely, u_j is given by

$$(2.14) \quad (2\pi)^{-k} t^{m-k_j/2+(2n+1)/4} x^{r-k/2+(2n-1)/4} \int_E e^{i\psi_j(x,y,w,v)/xt} a_j(x, y, w, v, t) dv.$$

Here $v \in \mathbb{R}^{k_j}$, $E \subset \mathbb{R}^{k_j}$ is compact, a_j is a smooth function of x, y, w, v with compact support and ψ is a nondegenerate parametrization of L on the support of a . Similarly, u_f

is given by a sum of terms of the form

$$(2.15) \quad (2\pi)^{-k} x^{r-k/2+(2n-1)/4} \int_E e^{i\psi_j(0,y,w,v)/xt} b_j(x, y, w, v, t) dv.$$

with b smooth and supported away from $t = 0$. It is straightforward to check that (2.14) and (2.15) are compatible. Note that, for fixed $t > 0$, the expression (2.15) is a Legendrian distribution of order $r - 1/4$ associated to the Legendre submanifold $t^{-1}G$, where t^{-1} acts by scaling in the fibre variables.

Fibred-scattering Legendrian distributions of order (m, r) on L have a well-defined symbol map σ^m taking values in $x^{r-m}\mathcal{C}^\infty(L; S^{[m]}(L))$ over L , which is defined by continuity from the symbol map in the interior of L . (This makes sense because the fibred-scattering structure at mf and away from sf is just the scattering structure.) The bundle $S^{[m]}(L)$ in the fibred-scattering setting is given by $S^{[m]}(L) = \Omega_b^{1/2}(L) \otimes |N^* \text{mf}|^{m-N/4} \otimes |N^* \text{sf}|^{m-N/4} \otimes M \otimes E$, where $\Omega_b^{1/2}(L)$ denotes the b -half-density bundle over L ; a smooth nonzero section near the boundary is $|d\lambda dx/x|^{1/2}$, where λ are coordinates on ∂L , extended into the interior of L . The symbol determines $u \in I^{m,r}(L)$ up to an element of $I^{m+1,r}(L)$.

3. GEOMETRY OF THE TIME-DEPENDENT SCHRÖDINGER OPERATOR

3.1. Flowout from the diagonal. Let $P = t^2(D_t + H)$. We consider this operator as an element of order $(2, 0)$ of the scattering calculus for the manifold with boundary $X^\circ \times X^\circ \times \mathbb{R}_{\geq 0}$. We shall use coordinates $z, w, t, \zeta, \eta, \tau$ in the scattering cotangent bundle, as in (2.2). The boundary symbol of P at $t = 0$ is

$$p(z, w, \zeta, \eta, \tau) = -\tau + \frac{1}{2}|\zeta|_g^2.$$

The Hamiltonian flow, rescaled by factor of t^{-1} , is given by

$$(3.1) \quad \begin{aligned} \dot{z} &= \frac{1}{2} \frac{\partial |\zeta|_g^2}{\partial \zeta} & \dot{\zeta} &= \zeta - \frac{1}{2} \frac{\partial |\zeta|_g^2}{\partial z} \\ \dot{w} &= 0 & \dot{\eta} &= \eta \\ \dot{t} &= t & \dot{\tau} &= |\zeta|_g^2. \end{aligned}$$

The free Schrödinger propagator $c_n t^{-n/2} e^{i|z-w|^2/2t}$ is a Legendrian distribution parametrized by the phase function $|z-w|^2/2$. We look for an analogue in the general case.

Lemma 3.1. *There exists a smooth Legendre submanifold L of ${}^{\text{sc}}T_{\{t=0\}}^* X^\circ \times X^\circ \times \mathbb{R}_{\geq 0}$ contained in $\{p=0\}$ that is parametrized by the phase function $\Phi(z, w) = d(z, w)^2/2$ near the diagonal $z = w$.*

Proof. Since $p(z, w, d_z \Phi, d_w \Phi, \Phi) = 0$, the function Φ parametrizes a Legendrian contained in $\{p=0\}$ in the region where Φ is smooth, that is, within the injectivity radius. We define L to be the flowout along bicharacteristics from this region. L is given as a set by

$$(3.2) \quad (z, \hat{\zeta}) = \exp_{sg/2}(w, \hat{\eta}_0), \quad |\eta| = |\zeta| = s, \quad \tau = \frac{s^2}{2}, \quad \hat{\eta} = -\hat{\eta}_0, \quad s \in [0, \infty), \quad (w, \hat{\eta}_0) \in S^* X^\circ.$$

Since the Hamilton vector field never vanishes for $s > 0$, and since $\tau \rightarrow \infty$ as $s \rightarrow \infty$, the entire flowout is smooth. \square

3.2. Behaviour of L near the boundary. We next analyze the behaviour of L near the boundary of X in the first factor. To do this, we introduce the Lie algebra of vector fields given by the $C^\infty(X \times X \times \mathbb{R}_{\geq 0})$ -span of the vector fields

$$t^2 x^2 \partial_t, tx^3 \partial_x, tx^2 \partial_y, tx^2 \partial_w.$$

There is a vector bundle of which these are smooth sections forming a basis at every point. Let B denote the dual of this bundle; then a basis of the smooth sections of B is given locally near the boundary of X by

$$\frac{dt}{t^2 x^2}, \frac{dx}{x^3 t}, \frac{dy_i}{x^2 t}, \frac{dw_j}{x^2 t}.$$

Hence an arbitrary $q \in B$ may be written³

$$(3.3) \quad q = \frac{\kappa}{x^2} d\left(\frac{1}{t}\right) + \frac{\lambda dx}{x^3 t} + \frac{\mu dy}{x^2 t} + \frac{\xi dw}{x^2 t}.$$

Then $x, y, w, t, \lambda, \mu, \xi, \kappa$ are smooth coordinates on B , up to the corner $t = x = 0$. We remark that the bundle B is a scaled cotangent bundle mixing scattering behaviour at the face $t = 0$ with quadratic scattering behaviour at $x = 0$.

There is a natural vector bundle isomorphism from ${}^s T^*(X^\circ \times X^\circ \times \mathbb{R}_{\geq 0})$ to B over X° given by

$$(3.4) \quad (z, w, t, \zeta, \eta, \tau) \rightarrow (z, w, t, x\zeta, x\eta, x^2\tau)$$

in terms of the coordinates given above. Thus, the map rescales the cotangent variables by a power of x as we approach $x = 0$ — linear rescaling for the spatial cotangent variables and quadratic for the temporal cotangent variable. This counteracts the growth of these variables as the boundary $x = 0$ is approached, since we see from (3.2) that ζ, η grow linearly and τ quadratically in s (which expected to be asymptotic to x^{-1} as $s \rightarrow \infty$). Thus, if we map L into B via the isomorphism (3.4), we can expect it to remain ‘bounded’ at $x = 0$. By abuse of notation, we denote the image of L under (3.4) also by L . The Legendrian L , now regarded as a submanifold of B , is generated by Hamiltonian flow of $x^2 P$ starting from the diagonal.

Owing to the special structure (1.1) of the scattering metric g , the boundary symbol of $x^2 P$ equals

$$(3.5) \quad -\kappa + \frac{1}{2}(\lambda^2 + h^{ij} \mu_i \mu_j)$$

where h^{ij} is a smooth function of x and y . This yields (rescaled) Hamilton equations in B

$$(3.6) \quad \begin{aligned} \dot{x} &= \lambda x & \dot{\lambda} &= \lambda + 2\kappa - 2h^{ij} \mu_i \mu_j - \frac{x}{2} \frac{\partial h^{ij}}{\partial x} \mu_i \mu_j \\ \dot{y}_i &= h^{ij} \mu_j & \dot{\mu}_i &= (1 + 2\lambda) \mu_i - \frac{1}{2} \partial_{y_i} h^{jk} \mu_j \mu_k \\ \dot{w} &= 0 & \dot{\xi} &= (1 + 2\lambda) \xi \\ \dot{t} &= t & \dot{\kappa} &= \lambda^2 + 2\lambda\kappa + h^{ij} \mu_i \mu_j. \end{aligned}$$

The next lemma justifies our hope that the Legendrian L stays bounded in B as we approach the boundary, $x \rightarrow 0$.

Lemma 3.2. *Along every non-trapped bicharacteristic ray inside the Legendrian L , we have $x \rightarrow 0$, $\xi \rightarrow 0$, $\lambda \rightarrow -1$, $\mu \rightarrow 0$.*

³The variable μ here is not the same as the μ from Sections 1 and 2.

Proof. Along every nontrapped ray, we have $x \rightarrow 0$. Hence, there are points along the ray where x is arbitrarily small and where $\dot{x} < 0$. Since $\dot{x} = \lambda x$, there are points along the ray where x is arbitrarily small and where $\lambda < 0$.

Now we look at the equation for λ . Since the ray lies inside the characteristic variety of x^2P , we can substitute $2\kappa = \lambda^2 + h^{ij}\mu_i\mu_j$ on the right hand side of (3.6). Hence

$$(3.7) \quad \dot{\lambda} = \lambda + \lambda^2 - h^{ij}\mu_i\mu_j - \frac{x}{2} \frac{\partial h^{ij}}{\partial x} \mu_i\mu_j.$$

Since $h^{ij}(x)$ is positive definite for small x , uniformly over ∂X (since ∂X is compact),

$$-h^{ij}\mu_i\mu_j - \frac{x}{2} \frac{\partial h^{ij}}{\partial x} \mu_i\mu_j \leq 0 \text{ for } x < \epsilon.$$

Thus, starting from a point on the ray where $x < \epsilon$ and $\lambda < 0$, we have

$$\dot{x} = \lambda x, \quad \dot{\lambda} \leq \lambda + \lambda^2 \text{ if } x < \epsilon.$$

These equations imply that λ remains negative and that x is decreasing from this point forwards on the ray. Moreover, since $\lambda + \lambda^2 < 0$ for $-1 < \lambda < 0$, we see that $\limsup \lambda \leq -1$. In particular, from some point on, $\lambda < -3/4$. This implies that $\xi \rightarrow 0$ along the ray.

Now consider the equation for μ . It implies that

$$(h^{ij}\mu_i\mu_j)' = 2(1 + 2\lambda)h^{ij}\mu_i\mu_j + x\lambda \frac{\partial h^{ij}}{\partial x} \mu_i\mu_j.$$

Since eventually $\lambda < -3/4$ and $x < \epsilon$, positivity of h^{ij} implies that the right hand side is $\leq -1/2 h^{ij}\mu_i\mu_j$, say, from some point on, so $\mu \rightarrow 0$. Returning to the equation (3.7) for λ , we see that $\mu \rightarrow 0$ implies that $\lambda \rightarrow -1$, since (3.7) with the μ terms removed has an attracting fixed point at $\lambda = -1$. \square

3.3. Smoothness at the boundary. Throughout this subsection, we assume that the metric g is nontrapping.

Let $\Sigma(P)$ be the zero set of the boundary symbol of x^2P on $B \cap \{t = 0\}$. We identify this set with ${}^{\text{qsc}}T^*(X \times X^\circ)$ by mapping

$$\frac{\kappa}{x^2} d\left(\frac{1}{t}\right) + \frac{\lambda dx}{x^3 t} + \frac{\mu dy}{x^2 t} + \frac{\xi dw}{x^2 t} \quad \text{to} \quad \frac{\lambda dx}{x^3} + \frac{\mu dy}{x^2} + \frac{\xi dw}{x^2}.$$

In terms of coordinates, we map $(x, y, w, \kappa, \lambda, \mu, \xi)$ to $(x, y, w, \lambda, \mu, \xi)$. This is an isomorphism on $\Sigma(P)$ since κ is given by $|(\lambda, \mu)|_{g_{x,y}}^2/2$ on $\Sigma(P)$. The image of L under this identification is then a *Lagrangian* in ${}^{\text{qsc}}T^*(X \times X^\circ)$ (which we shall denote \tilde{L}). We emphasize that this Lagrangian lies over the interior of $X \times X^\circ$, not at the boundary. In terms of parametrizations, if Φ/t is a local parametrization for L , then Φ is a local parametrization of \tilde{L} .

It turns out that \tilde{L} is not smooth at $x = 0$. To recover smoothness, we shall blow up the submanifold

$$(3.8) \quad S = \{x = 0, \lambda = -1, \mu = 0, \xi = 0\}$$

where \tilde{L} meets the boundary (by Lemma 3.2). We shall use Melrose's notation and terminology; thus, we denote the manifold with corners obtained by blowing up the submanifold S by

$$[{}^{\text{qsc}}T^*(X \times X^\circ); S];$$

explicitly, this is the manifold with corners obtained by removing S and replacing it with its inward-pointing spherical normal bundle, with a natural C^∞ structure (see [14]). This

inward-pointing spherical normal bundle becomes a new boundary hypersurface of the blown-up space which we refer to as the ‘front face’ of the blowup. The lift to this blown-up space of the original boundary hypersurface $\partial X \times X^\circ$ will be denoted sf .

Lemma 3.3. *Let (X, g) be a manifold with boundary, with a nontrapping scattering metric. Let $S \subset {}^{\text{qsc}}T^*(X \times X^\circ)$ be given by (3.8), where x is a boundary defining function such that $|dx/x^2|_g = 1$ at ∂X . Then*

$$(3.9) \quad X_{\text{tS}} \equiv [{}^{\text{qsc}}T^*(X \times X^\circ); S] \setminus \tilde{\text{sf}}$$

is an affine bundle over $X \times X^\circ$, isomorphic (but not naturally isomorphic) to the scattering cotangent bundle ${}^{\text{sc}}T^*(X \times X^\circ)$. The Lagrangian \tilde{L} lifts to the space (3.9) to be a smooth manifold with boundary, such that $\partial\tilde{L}$ is contained in the boundary of ${}^{\text{sc}}T^*(X \times X^\circ)$ and such that \tilde{L} is transversal to the boundary there.

Proof. Performing the blowup amounts to introducing the coordinates on X_{tS} in a neighbourhood of the front face. The induced map from X_{tS} to $X \times X^\circ$ is a fibration in which each fibre is diffeomorphic to \mathbb{R}^{2n} . To specify the affine structure we introduce the coordinates $\Lambda = (\lambda + 1)/x$, $M = \mu/x$, $\Xi = \xi/x$ which are smooth on X_{tS} up to $x = 0$. The affine structure is defined by deeming Λ, M and Ξ to be affine coordinates on each fibre. This is well defined since a smooth change of variables in $(x, y, w, \lambda, \mu, \xi)$ which is linear on each fibre of ${}^{\text{qsc}}T^*(X \times X^\circ)$ induces a smooth change of variables in $(x, y, w, \Lambda, M, \Xi)$ which is *affine* on each fibre. (Observe that a coordinate change such as $x \rightarrow x + \alpha x^2$ induces a translation $\Lambda \rightarrow \Lambda + \alpha + O(x)$, so the coordinate changes are not in general linear on the fibres.) In terms of the new coordinates, a point

$$q = \frac{\lambda dx}{x^3} + \frac{\mu \cdot dy}{x^2} + \frac{\xi \cdot dw}{x^2} \in {}^{\text{qsc}}T^*(X \times X^\circ)$$

becomes

$$(3.10) \quad q = -\frac{dx}{x^3} + \Lambda \frac{dx}{x^2} + M \cdot \frac{dy}{x} + \Xi \cdot \frac{dw}{x}.$$

Mapping q to $\Lambda dx/x^2 + M dy/x + \Xi dx/x$ defines an affine bundle isomorphism between the blown-up space X_{tS} defined in (3.9) and ${}^{\text{sc}}T^*(X \times X^\circ)$. It is not, however, a canonical isomorphism — this is clear, since the scattering cotangent bundle has a linear structure which X_{tS} lacks. We remark that if we performed the blowup at the zero section of the quadratic-scattering cotangent bundle, that is, at $\lambda = 0$ instead of $\lambda = -1$, then the resulting space as in (3.9) *would* be naturally isomorphic to the scattering cotangent bundle.

Next, we show that \tilde{L} has the regularity specified in the statement of the lemma. Our method of proof is to lift the vector field V in (3.6) to the space $[{}^{\text{qsc}}T^*(X \times X^\circ); S]$, and divide by a boundary defining function for the front face, obtaining a new vector field W . Clearly integral curves of V are the same as integral curves of W . We claim that W is a smooth vector field on $[{}^{\text{qsc}}T^*(X \times X^\circ); S]$ which is tangent to $\tilde{\text{sf}}$ but transverse to the front face. Consequently, every integral curve of W must meet the boundary of $[{}^{\text{qsc}}T^*(X \times X^\circ); S]$ in the interior of the front face. Moreover, the fact that W is smooth and nonvanishing at the front face implies that the flowout \tilde{L} is smooth up to the front face and meets it transversally. Hence to prove the lemma, it is sufficient to prove the claim above.

Proving the claim is simply a matter of computing the vector field (3.6) in various coordinate systems valid in different regions of the blown-up space $[{}^{\text{qsc}}T^*(X \times X^\circ); S]$. In a neighbourhood of any point in the interior of the front face, we may use coordinates Λ, M, Ξ

defined above. In terms of these coordinates, the vector field takes the form

$$(3.11) \quad V = (-x + \Lambda x^2)\partial_x + xh^{ij}M_j\partial_{y_i} + x(-h^{ij}M_iM_j + \frac{x}{2}\frac{\partial h^{ij}}{\partial x}M_iM_j)\partial_\Lambda \\ + x(\Lambda M_i - \frac{1}{2}\frac{\partial h^{jk}}{\partial y_i}M_jM_k)\partial_{M_i} + x\Lambda\xi \cdot \partial_\Xi.$$

Dividing by x , which is locally a boundary defining function for the front face in this region, shows that $W = V/x$ is a smooth vector field in this region which is transverse to the boundary.

Near the corner of $[\text{qsc}T^*(X \times X^\circ); S]$, i.e. near the boundary of the front face, we need several different coordinate patches. Let $\rho = \sqrt{x^2 + (\lambda + 1)^2 + |\xi|^2 + |\mu|^2}$, and let $s = \lambda + 1$. When $s \geq \rho/4$, then coordinates $s_1, r_1 = x/s_1, M_1 = \mu/s_1, \Xi_1 = \xi/s_1$ may be used. When $|\mu| \geq \rho/4$, then valid coordinates are $s_2 = |\mu|, r_2 = x/s_2, \Lambda_2 = (\lambda + 1)/s_2, \hat{\mu} = \mu/s_2, \Xi_2 = \xi/s_2$ may be used, and when $|\xi| \geq \rho/4$, then valid coordinates are $s_3 = |\xi|, r_3 = x/s_3, \Lambda_3 = (\lambda + 1)/s_3, M_3 = \mu/s_3, \hat{\xi} = \xi/s_3$. The union of these coordinate patches covers a neighbourhood of the corner of $[\text{qsc}T^*(X \times X^\circ); S]$.

In the first set of coordinates, (s_1, r_1, M_1, Ξ_1) , the vector field V takes the form (where the subscript 1 is dropped for notational convenience)

$$(3.12) \quad V = (-s + s^2k)\partial_s - rsk\partial_r + sh^{ij}M_j\partial_{y_i} \\ + s((1 - sk)M_i - \frac{1}{2}\frac{\partial h^{jk}}{\partial y_i}M_jM_k)\partial_{M_i} + s(1 - sk)\Xi_i\partial_{\Xi_i}$$

The vector field $W = V/s$ is tangent to $\tilde{\text{sf}}$ and transverse to the front face in the region of validity of these coordinates. The calculation for the second and third set of coordinates is similar, and is left to the reader. \square

Now let G be the boundary of \tilde{L} : $G = \partial\tilde{L} \subset \text{sc}T^*_{\partial X \times X^\circ}(X \times X^\circ)$.

Lemma 3.4. *The submanifold G of $\text{sc}T^*_{\partial X \times X^\circ}(X \times X^\circ)$ is Legendrian.*

Proof. The contact form on $\text{sc}T^*_{\partial X \times X^\circ}(X \times X^\circ)$ is given by $\omega(x^2\partial_x, \cdot)$, where the symplectic form ω is now given by

$$(3.13) \quad \omega = \frac{d\Lambda \wedge dx}{x^2} + \frac{dM \wedge dy}{x} + \frac{d\Xi \wedge dw}{x} - \frac{Mdx \wedge dy}{x^2} - \frac{\Xi dx \wedge dw}{x^2}$$

which we obtain by taking the differential of (3.10). Since V in (3.11) is of the form $-x\partial_x + xV''$, where V'' is tangent to the boundary of $\text{sc}T^*(X \times X^\circ)$, and $x\omega$ by (3.13) is nondegenerate on vector fields tangent to the boundary, uniformly up to the boundary, we find that the contraction of ω and $x^2\partial_x$ satisfies

$$\omega(x^2\partial_x, \cdot) = -x\omega(V, \cdot) + x\alpha$$

where α is a smooth one-form on $\text{sc}T^*(X \times X^\circ)$. Let W be a vector tangent to G . Then

$$\omega(x^2\partial_x, W) = -x\omega(V, W) + x\alpha(W),$$

The first term vanishes since $G \subset \tilde{L}$, both V and W are tangent to \tilde{L} , and \tilde{L} is Lagrangian. The second term vanishes at $x = 0$. This proves that G is Legendrian. \square

3.4. Asymptotics of geodesic flow and the Legendrian G . The variables Λ and M have geometric meaning on the Legendrian G in terms of the asymptotic behaviour of geodesics. Notice that, by (3.2), the value of τ on the Legendrian L is given by $d(z, w)^2/2$. Thus, κ is given by

$$\kappa = x^2 \tau = x^2 \frac{d(z, w)^2}{2}.$$

Near the boundary G of the Legendrian, we have by (3.5)

$$\kappa = \frac{1}{2}(\lambda^2 + h^{ij} \mu_i \mu_j) = \frac{1}{2}((-1 + x\Lambda)^2 + x^2 h^{ij} M_i M_j).$$

This implies that

$$(3.14) \quad d(z, w) = \frac{1}{x} - \Lambda + O(x).$$

Thus we interpret $-\Lambda$ on G as a ‘sojourn time’, the time that the geodesic spends in the interior of the manifold before emerging into the conic region.

Remark. It is instructive to note that under a change of boundary defining function $x \rightarrow \tilde{x} = x + \alpha x^2$ which preserves the normal form (1.1) we find that the sojourn time changes as follows: letting t be a unit speed parameter along the geodesic flow, we compute

$$\lim t - \frac{1}{\tilde{x}} = \lim t - \frac{1}{x} \frac{x}{\tilde{x}} = -\alpha + \lim t - \frac{1}{x};$$

hence we easily verify that this ‘sojourn time,’ while ill-defined as a function, does transform as a coordinate on the affine bundle $X_{\text{ts}}|_{x=0}$, as we know it must.

To interpret the coordinate M , we notice that according to (3.11) we have

$$(3.15) \quad \begin{aligned} \dot{y}_i &= h^{ij} \mu_j = x h^{ij} M_j, \\ \dot{x} &= -x + O(x^2), \end{aligned} \quad \text{which implies } dy_i/dx = -h^{ij} M_j + O(x).$$

Hence M measures the *angle* at which the geodesic strikes the boundary, or more geometrically, M is a coordinate on the pencil of ‘asymptotically parallel’ rays with a fixed final direction y . The coordinate y , of course, is just the limiting direction of the geodesic, while w and $-\hat{\Xi}$, which are constant under the flow, give the initial conditions of the geodesic.

3.5. Fibred scattering structure. There is yet another way of viewing the Legendrian L , in terms of the fibred-scattering structure of X_t^2 described in Section 2.3. We show that one can map the Legendrian $L \subset B$ to a fibred-scattering Legendrian ${}^\phi L$ of ${}^\phi T^* X_t^2$. To define this map, we use local coordinates (x, y, w, t) . Let

$$(3.16) \quad q = \frac{\kappa}{x^2} d\left(\frac{1}{t}\right) + \frac{\lambda dx}{x^3 t} + \frac{\mu dy}{x^2 t} + \frac{\xi dw}{x^2 t},$$

be a point in B , using coordinates as in (3.3). We map $q \in L \subset B$ to

$$(3.17) \quad q \mapsto q + \frac{1}{2} \frac{dt}{x^2 t^2} + \frac{dx}{x^3 t} = q - \frac{1}{2} d\left(\frac{1}{x^2 t}\right) \in {}^\phi T^* X_t^2.$$

Lemma 3.5. *Assume that the metric g is nontrapping. Then the image of (3.17) is a smooth fibred-scattering Legendrian submanifold ${}^\phi L$ of ${}^\phi T^* X_t^2$.*

Remark. Notice that the terms $dt/t^2 x^2$ and $dx/x^3 t$ in (3.16) are too ‘big’ for the fibred scattering space ${}^\phi T^* X_t^2$. The addition of terms as in (3.17) cancels the big part of these terms and yields something which remains ‘bounded’ in ${}^\phi T^* X_t^2$ up to $x = 0$.

Proof. We use the blowup coordinates Λ, M, Ξ as before. Thus,

$$\begin{aligned}\lambda &= -1 + x\Lambda \\ \mu &= xM \\ \xi &= x\Xi \\ \kappa &= \frac{1}{2}(\lambda^2 + h^{ij}\mu_i\mu_j) = -\frac{1}{2} - x\Lambda + x^2(\Lambda^2 + h^{ij}M_iM_j)\end{aligned}$$

Now we add $\frac{1}{2}dt/x^2t^2 + dx/x^3t$ to q in (3.16), and get

$$q \mapsto \Lambda \frac{xdt + tdx}{x^2t^2} + (\Lambda^2 + h^{ij}M_iM_j) \frac{dt}{t^2} + M \frac{dy}{xt} + \Xi \frac{dw}{xt}.$$

Thus there is a diffeomorphism between \tilde{L} and ${}^\phi L$, given in local coordinates by

$$(x, y, w, \Lambda, M, \Xi) \mapsto (x, y, w, \Lambda, \Lambda^2 + h^{ij}M_iM_j, M, \Xi),$$

so ${}^\phi L$ is smooth up to the boundary and meets it transversally. The submanifold ${}^\phi L$ is a Legendrian distribution. Indeed, if Ψ locally parametrizes L away from the boundary of B , then $\Psi - 1/(2x^2t)$ locally parametrizes ${}^\phi L$. Finally, under our identifications, the image of the boundary of ${}^\phi L$ under β is identical with G from Lemma 3.4, which is a Legendrian submanifold. This proves that ${}^\phi L$ is a fibred-scattering Legendre submanifold. \square

3.6. Trapped rays. If the metric g has trapped rays, then we need to localize the constructions of this section in the nontrapping region. We define L to be as in Lemma 3.1, but taking only those $(w, \hat{\eta}) \in \mathbf{F}$, rather than all $(w, \hat{\eta}) \in S^*X^\circ$. (Recall that \mathbf{F} denotes the set of points in S^*X° that are not trapped under the forward geodesic flow.) We then define $\tilde{L}, {}^\phi L$ and G in terms of L as before. Then Lemmas 3.3, 3.4 and 3.5 continue to hold except we lose smoothness of L, \tilde{L} and ${}^\phi L$ at the set corresponding to $s = 0$ in (3.2). Moreover, if we localize to a compact subset of \mathbf{F} , then the corresponding pieces of L, \tilde{L} and ${}^\phi L$ are compact.

Lemma 3.6. *The map S_f defined by*

$$(3.18) \quad S_f(w, \hat{\Xi}) = (y, \Lambda, M) \iff (y, w, \Lambda, M, -\Xi) \in G, \quad \frac{\Xi}{|\Xi|} = \hat{\Xi}$$

*is a contact diffeomorphism $S^*X^\circ \supset \mathbf{F} \mapsto {}^{\text{sc}}T_{\partial X}^*(X)$ satisfying (1.4) and (1.5). Similarly, $S_b(w, \hat{\Xi}) = -S_f(w, -\hat{\Xi})$ is a contact diffeomorphism.*

Proof. For each $(w, \hat{\Xi}) \in \mathbf{F}$ there is a unique point $(y, w, -\Lambda, -M, \Xi) \in G$ found by following the bicharacteristic emanating from $(w, \hat{\Xi})$ until it hits the front face of X_{tS} ; uniqueness and smoothness are a consequence of the transversality of the vector field W in the proof of Lemma 3.3. Invertibility of the map S_f follows from the previous discussion. In fact, given $(y, \Lambda, M) \in {}^{\text{sc}}T_{\partial X}^*(X)$, y determines a pencil of geodesics with asymptotic direction y , M picks out a unique geodesic within the pencil and then Λ indicates how far along the geodesic one must travel to get to the initial point w . The direction Ξ at w is the direction of that geodesic at w . Hence for each (y, Λ, M) there is a unique $(w, \hat{\Xi})$ with $S_f(w, \hat{\Xi}) = (y, \Lambda, M)$. Comparison between (2.1) and (3.10) shows that $\Lambda = -\nu$ when interpreted as a coordinate on ${}^{\text{sc}}T^*X$, hence (3.14) implies (1.4), while (3.15) implies (1.5).

The fact that S_f is contact follows directly from the definition (3.18) and the fact that G is Legendrian (see [6], p149, for a discussion in the symplectic case). \square

4. PROPAGATOR (NONTRAPPING CASE)

In this section, we shall construct a parametrix $\mathcal{U}(z, w, t)$ for the Schrödinger propagator, where we restrict the second variable w to some arbitrary open set \mathcal{G} with compact closure in X° . We require that \mathcal{U} solves

$$(4.1) \quad P\mathcal{U}(z, w, t) \equiv t^2(D_t + H_z)\mathcal{U}(z, w, t) = \mathcal{E}(z, w, t) \in t^N x^N \mathcal{C}^\infty(X \times \mathcal{G} \times \mathbb{R}_{\geq 0}) \text{ for all } N,$$

or in other words, the error term \mathcal{E} is smooth and rapidly decreasing both as $t \rightarrow 0$ and as the z variable tends to infinity. The restriction to $t > 0$ is only for convenience. In addition \mathcal{U} should satisfy the initial condition

$$(4.2) \quad \mathcal{U}(\cdot, w, t) \rightarrow \delta_w \text{ in } C^{-\infty}(X) \text{ as } t \rightarrow 0, \text{ for all } w \in \mathcal{G}.$$

In this section, we deal with the case in which the metric is globally nontrapping; we sketch the changes necessary to localize the construction in \mathbf{F} in the following section.

The parametrix will be a sum of four terms, $\mathcal{U} = \mathcal{U}_1 + \mathcal{U}_2 + \mathcal{U}_3 + \mathcal{U}_4$. Correspondingly, the construction is divided into four steps. We first construct a \mathcal{U}_1 supported near $t = 0$ and near the diagonal, satisfying the initial condition (4.2) and solving

$$(4.3) \quad P\mathcal{U}_1 = \mathcal{E}_1 + \mathcal{R}_1,$$

where \mathcal{E}_1 satisfies the condition for the error term in (4.1), and $\mathcal{R}_1(z, w, t)$ is supported in $\iota(X, g)/4 \leq d(z, w) \leq \iota(X, g)/2$, where ι is the injectivity radius of (X, g) . In the second step, we solve

$$(4.4) \quad P\mathcal{U}_2 = -\mathcal{R}_1 + \mathcal{E}_2 + \mathcal{R}_2,$$

with zero initial conditions, where \mathcal{E}_2 satisfies the condition for the error term in (4.1), and with $\mathcal{R}_2(z, w, t)$ supported in $d(z, w) \geq \iota(X, g)/4$, $t \leq 1$ and with $x \geq 2\epsilon$. (Here, and henceforth, we use (x, y) as local coordinates for the z variable when it is close to the boundary.) In the third step, we solve

$$(4.5) \quad P\mathcal{U}_3 = -\mathcal{R}_2 + \mathcal{E}_3 + \mathcal{R}_3,$$

with zero initial conditions, where \mathcal{E}_3 satisfies the condition for the error term in (4.1), and with $\mathcal{R}_3(z, w, t)$ supported in $x \leq 4\epsilon$. Finally, we solve

$$(4.6) \quad P\mathcal{U}_4 = -\mathcal{R}_3 + \mathcal{E}_4,$$

with zero initial conditions, where \mathcal{E}_4 satisfies the condition for the error term in (4.1). Clearly the sum $\mathcal{U}_1 + \mathcal{U}_2 + \mathcal{U}_3 + \mathcal{U}_4$ satisfies the conditions for a parametrix.

4.1. Step 1 — near the diagonal. We start by constructing a formal solution to

$$P\tilde{\mathcal{U}}_1(z, w, t) = t^2(D_t + H)\tilde{\mathcal{U}}_1(z, w, t) = 0, \quad t > 0,$$

near $z = w$, with initial condition a delta function $\delta_w(z)$. Our ansatz is

$$(4.7) \quad \tilde{\mathcal{U}}_1 = t^{-n/2} e^{i\Phi(z, w)/t} \sum_{j=0}^{\infty} t^j a_j(z, w).$$

By a formal solution we mean that each coefficient of t^j vanishes (near $z = w$) after applying the operator. Applying the operator, we find

$$(4.8) \quad \begin{aligned} t^{n/2} P\tilde{\mathcal{U}}_1 &= e^{i\Phi(z,w)/t} (-\Phi + g(\nabla_z \Phi, \nabla_z \Phi)) \sum_{j=0}^{\infty} t^j a_j \\ &+ it \left(-g(\nabla \Phi, \nabla) a_0 + \frac{n}{2} a_0 + \frac{1}{2} (\Delta \Phi) a_0 \right) \\ &+ \sum_{j=1}^{\infty} it^{j+1} \left(-g(\nabla \Phi, \nabla) a_j + \left(\frac{n}{2} - j \right) a_j + \frac{1}{2} (\Delta \Phi) a_j + \frac{i}{2} \Delta a_{j-1} - iV a_{j-1} \right) \end{aligned}$$

This gives us a sequence of equations to be solved, one for each power of t . The first equation is the *eikonal equation*, $-\Phi + g(\nabla_z \Phi, \nabla_z \Phi) = 0$. This has an exact solution

$$(4.9) \quad \Phi = \frac{1}{2} d(z, w)^2,$$

which is smooth when $d(z, w)$ is smaller than the injectivity radius of (X, g) . Motivated by the form of the free propagator on \mathbb{R}^n , we let this be our Φ .

The coefficients of t^j in the remainder are successively *transport equations* for a_0, a_1 , and so on. Consider the transport equation for a_0 . Fix a w and choose normal coordinates for z centred at $z = w$. Then

$$\Delta = \sum_j D_{z_j}^2 + O(z) D_z \text{ and } \Phi(z, w) = |z|^2/2 + O(|z|^3),$$

so

$$\Delta_z \Phi(z, w) + n = O(z) \text{ and } g(\nabla \Phi, \nabla) = \sum_j (z_j + O(|z|^2)) \partial_{z_j}.$$

Thus, the transport equation for a_0 has the form

$$(z_i + O(|z|^2)) \frac{\partial}{\partial z_i} a_0 = f \cdot a_0, \quad f = \frac{1}{2} \Delta \Phi + \frac{n}{2} = O(z),$$

where all terms are smooth. This has a unique smooth solution near the diagonal satisfying

$$(4.10) \quad a_0(z, z) = 1 \text{ for all } z \in X^\circ.$$

By the stationary phase lemma (see for example [10], Theorem 7.7.5), if Φ and a_0 satisfy (4.9) and (4.10) and all a_j are supported within half the injectivity radius of the diagonal, the initial condition (4.2) is satisfied.

The other transport equations take the form

$$(z_i + O(|z|^2)) \frac{\partial}{\partial z_i} a_j + j a_j = f \cdot a_j - \frac{i}{2} \Delta_z a_{j-1} - iV a_{j-1}, \quad f \text{ as above.}$$

We inductively suppose that a_0, \dots, a_{j-1} are smooth. Then there is a unique smooth solution a_j of this equation, establishing the inductive hypothesis for a_j .

To define \mathcal{U}_1 , we take our formal solution $\tilde{\mathcal{U}}_1$, and multiply by a smooth function $\chi(z, w)$ which is equal to 1 when $d(z, w) \leq \iota(X, g)/4$, and equal to 0 when $d(z, w) \geq \iota(X, g)/2$. Then we take \mathcal{U}_1 to be an asymptotic sum of the formal series so obtained. The error term we decompose into $\mathcal{E}_1 + \mathcal{R}_1$, where \mathcal{R}_1 is given by all terms containing a derivative of χ , and \mathcal{E}_1 is the remainder. Since $\tilde{\mathcal{U}}_1$ is a formal solution, \mathcal{E}_1 is $O(t^\infty)$ as $t \rightarrow 0$, while \mathcal{R}_1 is supported away from the diagonal. This completes Step 1.

4.2. Step 2 — at the main face. Here we solve away the error term \mathcal{R}_1 . We regard \mathcal{R}_1 as a half-density by multiplication by the Riemannian half-density $|dg_z dg_w dt|^{1/2}$. Then \mathcal{R}_1 has an asymptotic expansion of the form

$$(4.11) \quad t^{-n/2+1} e^{i\Phi(z,w)/t} \sum_{j=0}^{\infty} t^j r_j(z,w) |dg_z dg_w dt|^{1/2}.$$

It is therefore a Legendrian distribution, associated to the Legendrian L of Lemma 3.1 — see [15], or section 2.1 for a brief description.

We now need to specify how our differential operator P acts on half-densities; we shall do this by specifying a flat (i.e. covariant constant) half-density. One natural choice would be to specify that the Riemannian half-density $|dg_z dg_w dt|^{1/2}$ is flat. However, we shall make a different choice here since we want to apply the symbol calculus from [7], where it is assumed that the flat half-density is a scattering density (that is, a bounded non-vanishing section of the scattering half-density bundle). Thus, we specify instead that

$$(4.12) \quad \left| \frac{dg_z dg_w dt}{t^{2n+2}} \right|^{1/2} \text{ is flat, i.e. } \nabla \left| \frac{dg_z dg_w dt}{t^{2n+2}} \right|^{1/2} = 0.$$

For convenience we define

$$(4.13) \quad \alpha = \left| \frac{dg_z dg_w dt}{t^{2n+2}} \right|^{1/2}.$$

Hence, we want to solve

$$(4.14) \quad P(\mathcal{R}_2 \alpha) = -\mathcal{R}_1 \alpha,$$

or equivalently

$$(4.15) \quad (P + i(n+1)t)(\mathcal{R}_2 |dg_z dg_w dt|^{1/2}) = -\mathcal{R}_1 |dg_z dg_w dt|^{1/2}.$$

The additional term $i(n+1)t$ is a subprincipal term, since it vanishes to first order at $t = 0$.

Writing \mathcal{R}_1 in terms of α and using (2.14) we see that $\mathcal{R}_1 \in I^{7/4}(L)$. The Legendrian L is characteristic for P (that is, the symbol p of P vanishes on L , by construction of L), so we look for $\mathcal{U}_2 \in I^{3/4}(L; {}^{\text{sc}}\Omega^{1/2})$ whose symbol satisfies the transport equation (2.7) along L . In our situation, ${}^{\text{sc}}H_p$ is given by (3.1), while $\partial p / \partial \tau = 1$, and the subprincipal symbol $p_{\text{sub}} = i$ according to formula (2.9) of [7]. Therefore, by (3.1), we can solve away the error term $\mathcal{R}_1 \in I^{7/4}(L)$ with a Legendrian distribution u_0 of order $3/4$, by solving the ODE

$$(4.16) \quad -i \left(\mathcal{L}_{H_p} - \frac{n}{2} \right) \sigma^{3/4}(u_0) = \sigma^{7/4}(\mathcal{R}_1)$$

with ‘initial condition’ that the symbol of u_0 vanishes in a neighbourhood of $\{z = w, \zeta = \eta = 0\}$. All bicharacteristics originate here and tend to the boundary $\{x = 0\}$ by the non-trapping assumption, so there is a unique smooth solution with this property. It follows that in the region $x \geq \epsilon$, we have

$$Pu_0 - \mathcal{R}_1 \in I^{7/4+1}(L)$$

Inductively, assuming that $u_k \in I^{3/4}(L)$ solves

$$Pu_k - \mathcal{R}_1 \in I^{7/4+k+1}(L)$$

in the region $x \geq \epsilon$, we can find a $u_{k+1} \in I^{3/4}(L)$ which solves

$$Pu_{k+1} - \mathcal{R}_1 \in I^{7/4+k+2}(L)$$

in the region $x \geq \epsilon$, by solving

$$(4.17) \quad -i \left(\mathcal{L}_{H_p} + k - \frac{n}{2} \right) \sigma^{3/4+k+1}(v_k) = \sigma^{7/4+k+1}(Pu_k - \mathcal{R}_1)$$

and letting $u_{k+1} = u_k + v_k$. The v_k can then be asymptotically summed to yield a $u \in I^{3/4}(L)$ which solves

$$Pu - \mathcal{R}_1 \in t^N \mathcal{C}^\infty(\{x \geq \epsilon\} \times \mathcal{G} \times \mathbb{R}_{\geq 0}) \text{ for all } N.$$

We choose a cutoff function $\chi_2 \in \mathcal{C}^\infty(X)$ which is equal to 1 when $x \geq 2\epsilon$ and 0 when $x \leq \epsilon$. Define

$$\mathcal{U}_2 = \chi_2 u,$$

then \mathcal{U}_2 satisfies

$$P\mathcal{U}_2 = -\mathcal{R}_1 + \mathcal{R}_2 + \mathcal{E}_2,$$

where \mathcal{R}_2 is a Legendre distribution of order $7/4$ supported in $\epsilon \leq x \leq 2\epsilon$ (the \mathcal{R}_2 error comes from derivatives hitting the cutoff function χ_2). Our new error term \mathcal{R}_2 is now localized near the corner.

4.3. Step 3 — near the corner. Recall that we expect the fundamental solution to have quadratic oscillations at spatial infinity, i.e. at $x = 0$. In fact, we expect that $e^{-i/2tx^2}U(t)$ is a somewhat simpler kernel than U itself, near infinity. Therefore, we look for \mathcal{U}_3 in the form

$$\mathcal{U}_3 |dg_z dg_w dt|^{1/2} = e^{i/2tx^2} \tilde{\mathcal{U}}_3 |dg_z dg_w dt|^{1/2},$$

where $\tilde{\mathcal{U}}_3$ solves

$$e^{-i/2tx^2} \left(P e^{i/2tx^2} (\tilde{\mathcal{U}}_3 \alpha) \right) = e^{-i/2tx^2} (-\mathcal{R}_2 \alpha).$$

Here α is the half-density (4.13). Then in terms of $\tilde{\mathcal{U}}_3$ we have

$$(4.18) \quad \left(t^2 D_t - tx D_x + t^2 H + it \frac{n+2}{2} - txq \right) \tilde{\mathcal{U}}_3 = -e^{-i/2tx^2} \mathcal{R}_2.$$

Here $q = D_x \log \sqrt{h}$ and the term $i(n+1)t$ comes from the fact that we have chosen the scattering half-density α rather than the Riemannian half-density to be covariantly constant, as in Step 2. Note that the differential operators appearing in (4.18) are generated by the fibred-scattering vector fields on X_t^2 appearing in (2.11). This is not too surprising, since the analytic operation of multiplying \mathcal{U}_3 by $e^{-i/2tx^2}$ corresponds to the geometric operation of (3.17) which maps the Lagrangian \tilde{L} to a smooth Legendrian ${}^\phi L$ in ${}^\phi T^* X_t^2$. Since ${}^\phi L$ is well-behaved in ${}^\phi T^* X_t^2$, in the sense of being smooth up to the boundary, we can expect the corresponding operator (4.18) to be well-behaved in terms of the fibred-scattering structure on ${}^\phi T^* X_t^2$.

Lemma 4.1. *The kernel $e^{-i/2tx^2} \mathcal{R}_2$ is an element of $I^{7/4, \infty}({}^\phi L)$, as defined in [7].*

Proof. If Ψ parametrizes the Legendrian L , then $\Psi - 1/(2tx^2)$ parametrizes ${}^\phi L$ according to Lemma 3.5. Hence $e^{-i/2tx^2} \mathcal{R}_2$ is Legendrian with respect to ${}^\phi L$, of order $7/4$ at mf according to the calculation in Step 2. Since \mathcal{R}_2 vanishes in a neighbourhood of $x = 0$, it is order ∞ at sf. \square

Thus, it makes sense to seek the solution $\tilde{\mathcal{U}}_3$ to (4.18) in the space $I^{3/4, *}({}^\phi L)$. Here the value of $*$, specifying the rate of decay of the symbol as $x \rightarrow 0$, is determined by the transport equation which is regular singular at $x = 0$.

To do this, we must analyze the Hamilton vector field of our operator on ${}^\phi L$ near the boundary at $x = 0$. This is given (modulo some identifications) by the vector field V in (3.11), which is $x\partial_x + xW$, where W is tangent to the boundary. To determine the subprincipal symbol, observe that the adjoint of $t(xD_x - in/2)$ with respect to the Riemannian density is $t(xD_x - in/2) + O(tx)$ near the corner, hence the Weyl symbol of $t(xD_x - in/2)$ is $\nu + O(tx)$ where ν is the variable dual to x . Hence the subprincipal symbol of $t(xD_x - in/2)$ at mf vanishes at the boundary of mf (that is, at $x = 0$). Also, the subprincipal symbol of t^2H vanishes identically on the main face. So the boundary subprincipal symbol of the operator in (4.18) is equal to $i + O(x)$. Thus, we look for a $u_0 \in I^{3/4,*}({}^\phi L)$ satisfying

$$(4.19) \quad \left(-i\mathcal{L}_{-x\partial_x+xW} - i\left(\frac{1}{2} + \frac{3}{4} - \frac{2n+1}{4}\right) + i + x\tilde{q} \right) \sigma^{3/4}(u_0) = \sigma^{7/4}(e^{-i/2tx^2} \mathcal{R}_2)$$

This gives an equation of the form

$$\left(\mathcal{L}_{x\partial_x+xW} + \frac{n}{2} + x\tilde{q} \right) \sigma^{3/4}(u_0) = -i\sigma^{7/4}(e^{-i/2tx^2} \mathcal{R}_2),$$

so by (2.6), $\sigma^{3/4}(u_0)$ is of the form $a_0 x^{-n/2} |x^{-1} dx d\lambda|^{1/2} |dt|^{3/4-(2n+1)/4}$, where a_0 is smooth, and λ are coordinates on the boundary of ${}^\phi L$, extended into the interior. Here we have explicitly included the power of $|dt|$ in the formula since we now need to change boundary defining function from t to $\rho = tx$, in order to apply formulae from [7]. Doing this, we find that $\sigma^{3/4}(u_0)$ is of the form $a_0 x^{-1/2} |x^{-1} dx d\lambda|^{1/2} |d\rho|^{3/4-(2n+1)/4}$, where a_0 is smooth. By the (corrected) symbol calculus for fibred Legendrians given by Proposition 3.4 of [7]⁴, this implies that $u_0 \in I^{3/4,1/4}({}^\phi L)$, and that

$$(4.20) \quad \left(t^2 D_t - tx D_x + t^2 H + it \frac{n+2}{2} - txq \right) u_0 + e^{-i/2tx^2} \mathcal{R}_2 \in I^{7/4+1,1/4}({}^\phi L).$$

Inductively, we look for $u_l \in I^{3/4,1/4}({}^\phi L)$ such that

$$(4.21) \quad \left(t^2 D_t - tx D_x + t^2 H + it \frac{n+2}{2} - txq \right) u_l + e^{-i/2tx^2} \mathcal{R}_2 \in I^{7/4+l+1,1/4}({}^\phi L).$$

In fact, we will show more — we will show that there is u_l as above whose (full) symbol is of the form

$$(4.22) \quad a(\rho, x) x^{-1/2} |x^{-1} dx d\lambda|^{1/2} |d\rho|^{3/4-(2n+1)/4}, \quad \text{with } a \text{ smooth.}$$

The important point here is that a is a smooth function of ρ and x , not just a smooth function of t and x . To show this, we use the boundary defining function ρ rather than t for the main face, even though it is degenerate at the corner $t = x = 0$. In terms of the coordinates ρ, x, y, w near the corner, partial derivatives transform as

$$(4.23) \quad \begin{aligned} tD_t|_{x,y} &= \rho D_\rho|_{x,y} \\ xD_y|_{x,t} &= xD_y|_{\rho,x} \\ xD_x|_{y,t} &= xD_x|_{y,\rho} + \rho D_\rho|_{x,y}. \end{aligned}$$

We also write $V = x^2 \tilde{V}$ where $\tilde{V} \in \mathcal{C}^\infty(X)$ by assumption. Hence the operator in (4.21) takes the form

$$(4.24) \quad \tilde{P} = -\rho D_x + \frac{\rho^2}{2} \left((xD_x + \rho D_\rho)^2 + i(n-2)(xD_x + \rho D_\rho) + \Delta_{h(x)} + \tilde{V} \right) + i\rho \frac{n+2}{2x} - \rho q.$$

⁴The power of ρ in the final nonzero term of the exact sequence of Proposition 3.4 of [7] is incorrect. It should be $\mathbf{r} - m$, not $m - \mathbf{r}$.

Any Legendrian distribution u in $I^{3/4,1/4}(\phi L)$ can be written, modulo $x^\infty t^\infty C^\infty(X \times \mathbb{R}_{\geq 0})$, as a finite sum of oscillatory integrals of the form (written as a Taylor series in ρ)

$$(4.25) \quad u = \int_K e^{i\psi(x,y,w,v)/\rho} \rho^{n/2+1-k/2} \sum_{j=0}^{\infty} \rho^j a_j(x, y, w, v) dv$$

where ψ parametrizes ϕL , $K \subset \mathbb{R}^k$ is bounded, and $a_j \in x^{-n/2-1-j} C^\infty$. We claim that in fact, $a_j \in x^{-n/2-1} C^\infty$, or in other words, that the full symbol of u is smooth in x and ρ .

The equation to be solved is

$$(4.26) \quad \tilde{P}u = -e^{-i/2tx^2} \mathcal{R}_2 = \int_K e^{i\psi(x,y,w,v)/\rho} \rho^{n/2+2-k/2} \sum_{j=0}^{\infty} \rho^j b_j(x, y, w, v) dv$$

where the b_j are supported in $x \geq \epsilon$. Substituting (4.24) for \tilde{P} , we find that the left hand side is given by

$$(4.27) \quad \int_K e^{i\psi(x,y,w,v)/\rho} \rho^{n/2+1-k/2} \left\{ \left(-\psi_x + \frac{1}{2}(x\psi_x - \psi)^2 + \frac{1}{2}|d_y\psi|_{h(x)}^2 \right) + \rho \left(-D_x + (x\psi_x - \psi)(xD_x + \rho D_\rho) + \langle \nabla_y \psi, D_y \rangle_{h(x)} + i\frac{n+2}{2x} + f \right) + \rho^2 Q \right\} \sum_{j=0}^{\infty} \rho^j a_j(x, y, w, v) dv,$$

where Q is a second order operator generated over $\mathcal{C}^\infty(x, y)$ by $x D_x$, ρD_ρ , and D_y and $f \in \mathcal{C}^\infty(x, y)$. Since ψ parametrizes a Legendrian which is characteristic for \tilde{P} , the expression

$$(4.28) \quad -\psi_x + \frac{1}{2}(x\psi_x - \psi)^2 + \frac{1}{2}|d_y\psi|_{h(x)}^2$$

vanishes when $d_v\psi = 0$. As ψ satisfies the nondegeneracy condition (2.13), the function (4.28) can be expressed

$$\sum_{l=1}^k a_l \frac{\partial \psi}{\partial v_l} \quad \text{with } a_l \text{ smooth.}$$

We then write $a_l(\partial_{v_i}\psi)e^{i\psi/\rho} = a_l \rho D_{v_i} e^{i\psi/\rho}$ and integrate by parts in v . These terms then become $O(\rho)$ terms. Let us write $W = (x\psi_x - \psi)x D_x + \langle \nabla_y \psi, D_y \rangle_{h(x)} + \sum_l a_l D_{v_l}$. The equation becomes

$$(4.29) \quad \sum_{j=0}^{\infty} \rho^j \left\{ \left(-D_x + W + i\frac{n+2}{2x} + f_j \right) + \rho Q_j \right\} a_j = \sum_{j=0}^{\infty} \rho^j b_j.$$

Here f_j are smooth and Q_j is a second order operator generated by $x D_x$, D_y and D_v with smooth coefficients. Thus, we need to solve

$$(4.30) \quad \begin{aligned} (-xD_x + xW + i\frac{n+2}{2} + xf_0)a_0 &= xb_0, \\ (-xD_x + xW + i\frac{n+2}{2} + xf_j)a_j &= xb_j + xQ_j a_{j-1}. \end{aligned}$$

Here b_j is supported in $x \in [\epsilon, 2\epsilon]$ and our initial condition for a_j is that it vanish for $x > 2\epsilon$. These are regular singular ODEs for the a_j . The first equation certainly has a solution which is in $x^{-n/2-1} C^\infty$. Assuming that a_j is in $x^{-n/2-1} C^\infty$, it follows from (4.30) that $a_{j+1} \in x^{-n/2-1} C^\infty$, since the right hand side of (4.30) is in $x^{-n/2} C^\infty$ and the indicial root of the regular singular operator on the left hand side is $-n/2 - 1$. Hence the claim follows

by induction. (Notice that this induction would not work if we did not make the stronger assumption (4.22) on the symbol of u .)

Thus, we let $\tilde{\mathcal{U}}_3$ be an asymptotic sum of the formal series (4.25) constructed above, supported say in $t \leq 1$. This is certainly in $I^{3/4,1/4}(\phi L)$ and solves the equation

$$(4.31) \quad \left(t^2 D_t - tx D_x + t^2 H + it \frac{n+2}{2} - txq \right) \tilde{\mathcal{U}}_3 = -e^{-i/2tx^2} \mathcal{R}_2 + \mathcal{E}_3 + \tilde{\mathcal{R}}_3,$$

where $\tilde{\mathcal{R}}_3 \in I^{\infty,1/4}(\phi L)$. This completes Step 3.

4.4. Step 4 — near spatial infinity. Now we have an error term $\tilde{\mathcal{R}}_3 \in I^{\infty,1/4}(\phi L)$, and we seek a solution to

$$(4.32) \quad \left(t^2 D_t - tx D_x + t^2 H + it \frac{n+2}{2} - txq \right) \tilde{\mathcal{U}}_4 = -\tilde{\mathcal{R}}_3 \in x^\infty t^\infty C^\infty(X_t^2).$$

We seek a solution $\tilde{\mathcal{U}}_4$ in the space $I^{\infty,1/4}(\phi L)$.

We can write $-\tilde{\mathcal{R}}_3$ as a finite sum of terms of the form

$$(4.33) \quad \int_K e^{i\psi(x,y,w,v)/\rho} \rho^{-k/2} \sum_{j=0}^{\infty} \rho^j b_j(t, y, w, v) dv$$

where we have expanded the symbol as a Taylor series in ρ . Here each b_j is $O(t^\infty)$ at $t = 0$. In terms of the coordinates (t, ρ, y) ,

$$tD_t|_{\rho,y} = tD_t|_{x,y} - xD_x|_{t,y},$$

so the operator in (4.32) takes the form

$$(4.34) \quad t^2 D_t|_{\rho} + it \frac{n+2}{2} + (\rho^2 D_\rho)^2 + h^{ij}(\rho D_{y_i})(\rho D_{y_j}) + \frac{\rho}{t} Q - \rho q$$

where Q is a second order differential operator generated over $C^\infty(x, y)$ by the vector fields $\rho^2 D_\rho$ and ρD_y . Hence, we wish to solve

$$(4.35) \quad \left(t^2 D_t|_{\rho} + it \frac{n+2}{2} + (\rho^2 D_\rho)^2 + h^{ij} \rho^2 D_{y_i} D_{y_j} + \frac{\rho Q}{t} - \rho q \right) \int_K e^{i\psi/\rho} \sum_{j=0}^{\infty} \rho^{j-\frac{k}{2}} a_j(t, y, w, v) dv \\ = \int_K e^{i\psi/\rho} \sum_{j=0}^{\infty} \rho^{j-\frac{k}{2}} b_j(t, y, w, v) dv, \quad \psi = \psi(x, y, w, v).$$

Equation (4.35) holds (as a formal series in ρ) provided we have

$$(4.36) \quad \left(t^2 D_t + it \frac{n+2}{2} + c(y, w, v) \right) a_0 = b_0, \\ \left(t^2 D_t + it \frac{n+2}{2} + c(y, w, v) \right) a_j = b_j + R_j(a_0, \dots, a_{j-1}),$$

where $c(y, w, v) = \psi(0, y, w, v)^2 + h^{ij} \psi_{y_i}(0, y, w, v) \psi_{y_j}(0, y, w, v)$ is independent of t . The operator R_j here is the coefficient of ρ^j in the expansion of Q and is such that $t^{j+1} R_j$ is a second order differential operator in y with coefficients smooth in (y, t) . We show inductively that there is a solution with each $a_j \in t^\infty C^\infty$. Indeed, the equation for a_0 is explicitly solved by

$$a_0(t) = t^{(n+2)/2} e^{ic/t} \int_0^t i e^{-ic/s} b_0(s) s^{-n/2+3} ds$$

which is in $t^\infty C^\infty$ since $b_0 \in t^\infty C^\infty$. Inductively assuming that a_0, \dots, a_{j-1} are in $t^\infty C^\infty$, then it follows from the formula above, with b_0 replaced by $b_j + R_j(a_0, \dots, a_{j-1})$ that $a_j \in t^\infty C^\infty$. Let \tilde{U}_4 be an asymptotic sum of the series on the left hand side of (4.35). Then \tilde{U}_4 solves (4.32) up to an $x^\infty t^\infty C^\infty$ error term. This completes the construction of the parametrix in the nontrapping case.

4.5. Exact solution. We have now shown that

$$(4.37) \quad (D_t + H)\mathcal{U}(x, y, w, t) = e(x, y, w, t) \in \dot{C}^\infty(\mathbb{R}_{\geq 0} \times X \times \mathcal{G}) = \dot{C}^\infty(\mathbb{R}_{\geq 0}; \dot{C}^\infty(X \times \mathcal{G})).$$

The exact propagator $U(t)$ is given in terms of the parametrix $\mathcal{U}(t)$ by Duhamel's formula

$$(4.38) \quad U(t) = \mathcal{U}(t) + i \int_0^t U(s)(D_t + H)\mathcal{U}(t-s) ds.$$

It follows from a commutator argument due to Craig, [2] Théorème 14, that, letting

$$\mathcal{H}_k = \bigcap_{s=0}^k x^{k-s} H_{sc}^s(X),$$

we have for all $k > 0$

$$(4.39) \quad U(t) : \mathcal{H}_k \rightarrow L_{loc}^\infty(\mathbb{R}_t; \mathcal{H}_k).$$

Note that $\bigcap_k \mathcal{H}_k = \dot{C}^\infty$, so in particular

$$(4.40) \quad U(t) : \dot{C}^\infty(X) \rightarrow \dot{C}^\infty(X) \quad \text{for all } t \leq 0.$$

Lemma 4.2. *Let $e \in \dot{C}^\infty(\mathbb{R}_{\geq 0}; \dot{C}^\infty(X \times \mathcal{G}))$. Then*

$$Ke(t)(z, w) \equiv \int_0^t ds \int_X U(s)(z, z')e(t-s)(z', w) dg(z') \in \dot{C}^\infty(\mathbb{R}_{\geq 0}; \dot{C}^\infty(X \times \mathcal{G})).$$

Proof. Let us fix $w \in \mathcal{G}$; then we may regard $e(z, w, t)$ as an element of $t^n \dot{C}^\infty(\mathbb{R}_{\geq 0}; \mathcal{H}_k)$ for any $n, k \in \mathbb{N}$. First we establish that $Ke \in t^n L^\infty(\mathbb{R}_{\geq 0}; \mathcal{H}_k)$. We compute for $t \in [0, T]$

$$\begin{aligned} \|Ke(t)\|_{\mathcal{H}_k} &\leq \int_0^t \|U(s)e(t-s)\|_{\mathcal{H}_k} ds \\ &\leq C_T \int_0^t \|e(t-s)\|_{\mathcal{H}_k} ds \\ &= O(t^n) \text{ for all } n \in \mathbb{N}. \end{aligned}$$

This shows that $Ke \in t^n L^\infty(t; \mathcal{H}_k)$ for every k and n . Derivatives of t and w can now be estimated similarly and uniform estimates then follow from the relative compactness of \mathcal{G} . \square

We conclude from (4.38) and Lemma 4.2 that $U(t) - \mathcal{U}(t) \in \dot{C}^\infty(\mathbb{R}_{\geq 0}; \dot{C}^\infty(X \times \mathcal{G}))$. This proves that the kernel of $U(t)$ is such that $e^{-i/2x^2 t} U(t) \in I^{3/4, 1/4}(\phi L)$, which completes the proof of Theorem 1.1.

5. THE TRAPPING CASE

We now sketch briefly the changes to be made in the parametrix construction in case there exist trapped geodesics. In this case, given a properly supported $Z \in \Psi^0(\mathcal{G})$ with $\text{WF}'Z \subset \mathbf{F}$, the non-forward-trapped part of the phase space, we wish to construct a partial parametrix \mathcal{U}_Z such that

$$(D_t + H)\mathcal{U}_Z(z, w, t) = e(z, w, t) \in \dot{\mathcal{C}}^\infty(\mathbb{R}_{\geq 0}; \dot{\mathcal{C}}^\infty(X \times \mathcal{G})).$$

with initial condition

$$\mathcal{U}_Z(t)f \rightarrow Zf \text{ in } C^{-\infty}(X) \text{ as } t \rightarrow 0.$$

It then follows by Duhamel's principle that $\mathcal{U}_Z - U(t)Z \in \dot{\mathcal{C}}^\infty(\mathbb{R}_{\geq 0}; \dot{\mathcal{C}}^\infty(X \times \mathcal{G}))$.

The only modification necessary in the construction comes at the end of Step 1. We begin with a result about the composition of pseudodifferential operators in the boundary with the simple Legendrian distributions appearing in Step 1.

Lemma 5.1. *Let $k(z, w, t) = t^s a(z, w, t)e^{i\Phi(z, w)/t}$ such that a and Φ are smooth, a is compactly supported and $d_w\Phi \neq 0$ on the support of a . Let $B \in \Psi^0(X)$ be a properly supported, classical pseudodifferential operator. Then*

$$v = \int_X k(z, w, t)B(w, w') dg(w)$$

has the form

$$v = t^s \tilde{a}(z, w, t)e^{i\Phi(z, w)/t}, \quad \tilde{a} \text{ smooth,}$$

with

$$\text{WF}_{\text{sc}}(v) \subset \text{WF}_{\text{sc}}(k) \cap \{(z, w, \zeta, \eta, \tau) \mid (w, \eta) \in \text{WF}'(B)\}.$$

Proof. The result follows directly from the lemma of stationary phase. Alternatively, we may regard z and t as smooth parameters and regard v as $B^T(k)$, the transpose of B acting on the distribution k . Recall that when B is a zeroth order classical pseudodifferential operator and $c, \phi \in \mathcal{C}^\infty$ with c compactly supported and $d\phi \neq 0$ on $\text{supp } c$,

$$B(c(w)e^{i\phi(w)/t}) = \tilde{c}(w, t)e^{i\phi(w)/t}$$

where $\tilde{c} \in \mathcal{C}^\infty(\mathbb{R}_{\geq 0} \times X)$ and $\tilde{c}(w, t) = O(t^\infty)$ for all w such that $(w, d_w\phi) \notin \text{WF}'B$. (See [20], Chapter 8, section 7.) The lemma now follows directly. \square

To construct \mathcal{U}_Z , we construct \mathcal{U}_1 as in Step 1 of the previous section and consider $\mathcal{U}_1 \circ Z$. The error term $\mathcal{R}_1 \circ Z = (D_t + H)\mathcal{U}_1 \circ Z$ is then a Legendre distribution associated to L , by the lemma just proved (since $d_w\Phi \neq 0$ on the support of the symbol of \mathcal{R}_1). Moreover the lemma shows that the scattering wavefront set of $(D_t + H)\mathcal{U}_1 \circ Z$ lies on the portion of L emanating from $(w, \eta) \in \text{WF}'(Z)$, which is contained in the non-forward trapped set. Now we proceed with Steps 2, 3 and 4 with $\mathcal{R}_1 \circ Z$ substituted for \mathcal{R}_1 . As no trapped rays lie in the microsupport of $\mathcal{R}_1 \circ Z$ however, we need only solve the transport equation in Step 2 along the remaining, non-trapped, rays. The rest of the construction proceeds as before.

Thus, we have shown that \mathcal{U}_Z is the sum of two terms, one of which has the form $\mathcal{U}_1 \circ Z$ where \mathcal{U}_1 is a Legendrian distribution as in subsection 4.1, and the other is a fibred-scattering Legendrian distribution of order $(\frac{3}{4}, \frac{1}{4})$.

Remark. It is not true that $\mathcal{U}_1 \circ Z$ is a Legendrian distribution associated to L , since $d_z\Phi = 0$ at $z = w$. In fact, $\mathcal{U}_1 \circ Z$ is a Legendrian conic pair associated to (Δ, L) , where Δ is the diagonal Legendrian $\{(z, \zeta, z, -\zeta, \kappa = 0)\}$. Since we do not need this fact, we omit the proof.

6. EGOROV THEOREM

Let $W(t) = e^{-i/2tx^2}U(t)$. Suppose A is a properly supported pseudodifferential operator on X° , microsupported in $O \times O$ with $O \subset \mathbf{B}$. We will show that $B = W(t)AW(t)^*$ is a scattering pseudodifferential operator of the same order as A , and that the symbol and microsupport of B are given by the pullback of the symbol and microsupport of A by the sojourn relation, scaled by the factor t^{-1} . Indeed we prove a more general result. To state it, suppose that S is a contact diffeomorphism from $\mathbf{O} \subset S^*X^\circ \rightarrow {}^{\text{sc}}T_{\partial X}^*X$. Then, by definition, $S^*\chi$, the pullback of the contact form on ${}^{\text{sc}}T_{\partial X}^*X$, is equal to a multiple $f\hat{\eta} \cdot dw$ of the contact form on S^*X . Associated to S is a Legendrian submanifold G of ${}^{\text{sc}}T_{\partial X \times X^\circ}^*X \times X^\circ$, given by

$$(6.1) \quad G = \{(y, w, \nu, \mu, -f\eta) \mid (w, \eta) \in \mathbf{O}, (y, \nu, \mu) = S(w, \eta)\}.$$

Indeed this is the correspondence of Lemma 3.6.

Proposition 6.1. *Let S be a contact diffeomorphism $S : \mathbf{O} \subset S^*X^\circ \rightarrow {}^{\text{sc}}T_{\partial X}^*X$ and let G be the Legendre submanifold given by (6.1). Suppose that $W \in I^0(X \times X^\circ, G; {}^{\text{sc}}\Omega^{\frac{1}{2}} \times \Omega^{\frac{1}{2}})$, and suppose that $A \in \Psi^m(X^\circ; \Omega^{\frac{1}{2}})$ is properly supported and microsupported in \mathbf{O} . Then*

$$(6.2) \quad WAW^* \in {}^{\text{sc}}\Psi^{-\infty, m}(X; {}^{\text{sc}}\Omega^{\frac{1}{2}}) \text{ with symbol } |\sigma(W)(q_G)|^2 \cdot \sigma(A)(S^{-1}(q))$$

where q_G is the point on G corresponding to q . Conversely, if $\tilde{A} \in {}^{\text{sc}}\Psi^{-\infty, m}(X; {}^{\text{sc}}\Omega^{\frac{1}{2}})$ has compact microsupport contained in the range of S , then modulo a kernel in $\dot{C}^\infty(X \times X)$, $W^*\tilde{A}W \in \Psi^m(X^\circ)$ is a properly supported pseudodifferential operator with symbol

$$(6.3) \quad \sigma(W^*\tilde{A}W)(\zeta) = \sigma(\tilde{A})(S(\zeta)) \cdot |\sigma(W)(\zeta_G)|^2.$$

The proof proceeds as follows: we reduce to the case $X = \overline{\mathbb{R}^n}$, the radial compactification of Euclidean space, by localization; applying the Fourier transform and results of Melrose-Zworski [15], we then deduce the result from the standard Egorov theorem.

To begin, we need a lemma, in effect a version of Proposition 10 of [15] with parameters, which tells us what happens when we pull back a Legendrian distribution on $\mathbb{R}^n \times \overline{\mathbb{R}^n}$ by the Fourier transform \mathcal{F} . The Fourier transform invariantly maps half-densities on a vector space to half-densities on the dual space. We recall from [15] that it can be interpreted as a Fourier integral operator associated to the Legendre diffeomorphism $L : {}^{\text{sc}}T_{\partial \overline{\mathbb{R}^n}}^*\overline{\mathbb{R}^n} \rightarrow S^*((\mathbb{R}^n)^*)$ given by $L(\hat{z}, \zeta) = (\zeta, -\hat{z})$, where $\hat{z} \in S^{n-1}$, the radial compactification of \mathbb{R}^n , and $\zeta \in (\mathbb{R}^n)^*$, the dual space.

Lemma 6.2. *Let S , G and W be as in Proposition 6.1 for $X = \overline{\mathbb{R}^n}$, and let \mathcal{F} be the Fourier transform. Then*

$$\mathcal{F} \circ W \in I^0((\mathbb{R}^n)^* \times \mathbb{R}^n, G_L; \Omega^{\frac{1}{2}}) + \mathcal{S}((\mathbb{R}^n)^* \times \mathbb{R}^n),$$

where G_L is the Lagrangian associated to the graph of the contact transformation $L \circ S$, L is the Legendre diffeomorphism, and \mathcal{S} denotes Schwartz space.

Proof of Lemma 6.2. Without loss of generality we may assume that W can be written in terms of a single parametrization. Then W can be written

$$(2\pi)^{-\frac{k}{2}} \int_{\mathbb{R}^k} e^{i\phi(\hat{z}, w, v)|z|} a(\hat{z}, \frac{1}{|z|}, w, v)|z|^{\frac{k}{2}} dv |dz dw|^{\frac{1}{2}}, \quad a \text{ smooth, compactly supported,}$$

modulo $\mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n)$, so $\mathcal{F} \circ W$ is given by

$$(2\pi)^{-n/2-k/2} \iint e^{-iz \cdot \zeta + i\phi(\hat{z}, w, v)|z|} a(\hat{z}, |z|^{-1}, w, v)|z|^{k/2} dv dz |d\zeta|^{\frac{1}{2}} |dw|^{\frac{1}{2}}$$

modulo $\mathcal{S}((\mathbb{R}^n)^* \times \mathbb{R}^n)$. Imitating the proof of Proposition 10 of [15], we set $\tilde{v} = v|z|$ so that $\mathcal{F} \circ W$ is given, modulo $\mathcal{S}((\mathbb{R}^n)^* \times \mathbb{R}^n)$, by

$$(6.4) \quad (2\pi)^{-n/2-k/2} \iint e^{-iz \cdot \zeta + i\phi(\hat{z}, w, \tilde{v}/|z|)|z|} a(\hat{z}, |z|^{-1}, w, \tilde{v}/|z|)|z|^{-k/2} d\tilde{v} dz |d\zeta|^{\frac{1}{2}} |dw|^{\frac{1}{2}}.$$

The phase is now homogeneous of degree one in (z, \tilde{v}) and parametrizes the Lagrangian G_L while the amplitude $a(\hat{z}, |z|^{-1}, w, \tilde{v}/|z|)|z|^{-k/2}$ has order $-k/2$ with $k+n$ phase variables z, \tilde{v} . Hence (6.4) is a Lagrangian distribution of Lagrangian order 0. \square

Proof of Proposition 6.1. By a microlocal partition of unity, we may assume that $\text{WF}' A \subset O$, a small subset of S^*X° . Then $S(O)$ lies over a small closed set in ∂X . Without loss of generality, we may assume that the support of the kernel $W(z, w)$ in the z variable lies within a small neighborhood W , covered by a single coordinate chart (x, y) , of this closed set, since introducing a spatial cutoff introduces only residual terms in WAW^* . By identifying W with a neighbourhood of a point of the boundary of $\overline{\mathbb{R}^n}$ and identifying a neighbourhood of $\pi(O)$ with a neighbourhood of the origin in \mathbb{R}^n , we may assume that that $A \in \Psi^0(\mathbb{R}^n)$ is properly supported and that W is as in Lemma 6.2.

Now we write

$$WAW^* = \mathcal{F}^*(\mathcal{F}W)A(\mathcal{F}W)^*\mathcal{F},$$

and apply the standard Egorov theorem to the middle three factors to conclude that this takes the form

$$\mathcal{F}^* \tilde{A} \mathcal{F},$$

where $\tilde{A} \in \Psi^m(\mathbb{R}^n) + \mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n)$ is, up to a Schwartz kernel, a pseudodifferential operator with principal symbol

$$\tilde{a}(q) = \sigma(A)(L^{-1}(S^{-1}(q))) \cdot \sigma((\mathcal{F}W)(\mathcal{F}W)^*)(q) = \sigma(A)(L^{-1}(S^{-1}(q))) \cdot |\sigma(W)(q_G)|^2.$$

Proposition 8 of [15] now shows that

$$\mathcal{F}^* \tilde{A} \mathcal{F} \in {}^{\text{sc}}\Psi^{-\infty, m}(\overline{\mathbb{R}^n})$$

with principal symbol

$$\sigma(A)(S^{-1}(q)) \cdot |\sigma(W)(q_G)|^2.$$

This proves (6.2). The proof of (6.3) is similar, proceeding from the expression

$$W^* \tilde{A} W = W^* \mathcal{F}^* \mathcal{F} \tilde{A} \mathcal{F}^* \mathcal{F} W.$$

\square

7. WAVEFRONT SET BOUND

In this section we prove Theorem 1.2. First we prove a preliminary result on the propagator at a fixed time $t \neq 0$. We define the contact transformations S_t for $t \neq 0$ as follows: for $t > 0$ and $q \in S^*X^\circ$, let $S_t(q)$ with domain \mathbf{F} be given by $S_t(q) = t^{-1}S_f(q)$ where S_f is the contact transformation defined in Lemma 3.6 and the scaling by t^{-1} acts in the fibre variables. For $t < 0$, let S_t with domain \mathbf{B} be given by $S_t(q) = |t|^{-1}S_b(q)$. Let G_t , $t \neq 0$, be the Legendrian (6.1) determined by S_t .

Lemma 7.1. *Let $Z \in \Psi^0(X^\circ)$ be properly supported, and microsupported inside \mathbf{F} . For fixed $t > 0$, the operator $W(t)Z$ is in $I^0(X \times X^\circ; G_t)$. Similarly, if $Z' \in \Psi^0(X^\circ)$ is properly supported, and microsupported inside \mathbf{B} , then for fixed $t < 0$, the operator $W(t)Z'$ is in $I^0(X \times X^\circ; G_t)$.*

Proof. The construction of the parametrix shows that $W(t)Z$ is given by a finite sum of oscillatory integrals

$$\int_{K \in \mathbb{R}^k} e^{i\psi(0,y,w,v)/xt} x^{-k/2} a(x,y,w,v,t) dv \left| \frac{dx dy dw}{x^{n+1}} \right|^{1/2},$$

where ψ and a are constructed in sections 4.3 and 4.4. (Here, because $t > 0$, we may replace $\psi(x,y,w,v)$ by $\psi(0,y,w,v)$ in the phase.) According to the remark below equation (2.15), this is a Legendre distribution associated to the Legendre submanifold G_t , of order zero. The proof for $t < 0$ is similar. \square

The proof of Theorem 1.2 is now straightforward.

Proof of Theorem 1.2. We only prove the theorem for $t > 0$, since the argument is similar for $t < 0$. Let $W(t) = e^{-i/2tx^2} U(t)$ as above. Then since both $U(t)$ and the multiplication operator $e^{-i/2tx^2}$ are unitary and map $\dot{C}^\infty(X)$ to itself, the same is true of $W(t)$. For simplicity we shall first assume that the metric is nontrapping.

Suppose that $q \notin \text{WF}(u(\cdot, t))$ for some $t > 0$. Then there is a properly supported $A \in \Psi^0(X)$ which is elliptic at q and such that $Au(\cdot, t) \in \dot{C}^\infty(X)$.

Since $f = U(-t)u$, we have $f = e^{-i/2tx^2} W(-t)u$, or $u = W(-t)^* e^{i/2tx^2} f$. Hence

$$Au = AW(-t)^* e^{i/2tx^2} f \in \dot{C}^\infty(X).$$

Applying $W(-t)$, we see that also $W(-t)AW(-t)^* e^{i/2tx^2} f \in \dot{C}^\infty(X)$. By Proposition 6.1, $W(-t)AW(-t)^* = \tilde{A}$ is a scattering pseudodifferential operator of order zero, which is elliptic at $S_{-t}(q)$. We have just shown that $\tilde{A}(e^{i/2tx^2} f) \in \dot{C}^\infty(X)$, so by definition of the scattering wavefront set, we see that $S_{-t}(q) \notin \text{WF}_{\text{sc}}(e^{i/2tx^2} f)$. Conversely, assume that $S_{-t}(q) \notin \text{WF}_{\text{sc}}(e^{i/2tx^2} f)$. Then there exists a scattering pseudodifferential operator \tilde{A} with compact microsupport which is elliptic at $S_{-t}(q)$ and such that $\tilde{A}(e^{i/2tx^2} f) \in \dot{C}^\infty(X)$. By Proposition 6.1, we may write $\tilde{A} = W(-t)AW(-t)^*$, with $A = A' + A'' \in \Psi^0(X^\circ) + \dot{C}^\infty(X \times X)$, where A' is properly supported and elliptic at q and A'' has a Schwartz kernel. Hence $W(-t)AW(-t)^* e^{i/2tx^2} f = W(-t)Au(\cdot, t)$ is in $\dot{C}^\infty(X)$, which implies (using (4.40)) that $Au(\cdot, t) \in \dot{C}^\infty(X)$. Hence $q \notin \text{WF}(u(\cdot, t))$.

Now we do not assume that the metric is nontrapping, but assume instead that the point q is not backward-trapped. Then since the set \mathbf{B} of non-backward-trapped points is open, one can choose A as above so that $\text{WF}'A \subset \mathbf{B}$. In fact, one can find in addition a properly supported pseudodifferential operator Z such that Z is equal to the identity microlocally on $\text{WF}'(A)$ and such that $\text{WF}'Z \subset \mathbf{B}$. Then $ZAZu(\cdot, t) \in \dot{C}^\infty(X)$. Repeating the argument above, we see that

$$W(-t)ZAZW(-t)^* e^{i/2tx^2} f \in \dot{C}^\infty(X).$$

In the trapping case, $W(-t)Z$ is a Legendre distribution associated to S_{-t} , so by Proposition 6.1, $W(-t)ZAZW(-t)^*$ is a scattering pseudodifferential operator which is elliptic at $S_{-t}(q)$, and the rest of the argument goes as before. \square

REFERENCES

- [1] N. Burq, P. Gérard and N. Tzvetkov, *Strichartz estimates and the nonlinear Schrödinger equation on compact manifolds*, Amer. J. Math., to appear.
- [2] W. Craig, *Les moments microlocaux et la régularité des solutions de l'équation de Schrödinger*, Séminaire sur les Équations aux Dérivées Partielles, 1995–1996, Sémin. Équ. Dériv. Partielles, École Polytech., Palaiseau, 1996, pp. Exp. No. XX, 24.

- [3] W. Craig, T. Kappeler, and W. Strauss, *Microlocal dispersive smoothing for the Schrödinger equation*, Comm. Pure Appl. Math. **48** (1995), no. 8, 769–860.
- [4] D. Fujiwara, *Remarks on convergence of the Feynman path integrals*, Duke Math. J. **47** (1980), no. 3, 559–600.
- [5] V. Guillemin, *Sojourn times and asymptotic properties of the scattering matrix*, Pub. RIMS **12**, supplement (1977), 69–88.
- [6] V. Guillemin and S. Sternberg, *Geometric Asymptotics*, revised edition, Mathematical surveys and monographs 14, American Mathematical Society, Providence RI.
- [7] A. Hassell and A. Vasy, *The resolvent for Laplace-type operators on asymptotically conic spaces*, Ann. Inst. Fourier (Grenoble) **51** (2001), no. 5, 1299–1346.
- [8] A. Hassell and A. Vasy, *The spectral projections and the resolvent for scattering metrics*, J. Anal. Math. **79** (1999), 241–298.
- [9] A. Hassell and J. Wunsch, *On the structure of the Schrödinger propagator*, to appear in *Partial Differential Equations and Inverse Problems*, Contemp. Math.
- [10] L. Hörmander, *The analysis of linear partial differential operators I*, Springer-Verlag, second edition, Berlin, 1990.
- [11] M. S. Joshi and A. Sá Barreto, *Recovering asymptotics of metrics from fixed energy scattering data*, Invent. Math. **137** (1999), no. 1, 127–143.
- [12] L. Kapitanski and Yu. Safarov, *A parametriz for the nonstationary Schrödinger equation*, Differential operators and spectral theory, Amer. Math. Soc. Transl. Ser. 2, vol. 189, Amer. Math. Soc., Providence, RI, 1999, pp. 139–148.
- [13] L. Kapitanski and Yu. Safarov, *Dispersive smoothing for Schrödinger equations*, Math. Res. Lett. **3** (1996), no. 1, 77–91.
- [14] R. B. Melrose, *Differential analysis on manifolds with corners*, manuscript in preparation, available on the web at <http://www-math.mit.edu/~rbm/book.html>.
- [15] R. B. Melrose and M. Zworski, *Scattering metrics and geodesic flow at infinity*, Invent. Math. **124** (1996), no. 1-3, 389–436.
- [16] R. B. Melrose, *Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces*, Spectral and scattering theory (Sanda, 1992), Dekker, New York, 1994, pp. 85–130.
- [17] L. Robbiano and C. Zuily, *Analytic theory for the quadratic scattering wave front set and application to the Schrödinger equation*, Univ. de Paris-Sud. Prépublication.
- [18] R. Schoen and S.-T. Yau, *Lectures on Differential Geometry*, International Press, 1994.
- [19] G. Staffilani and D. Tataru, *Strichartz estimates for a Schrödinger operator with nonsmooth coefficients*, Comm. PDE **27** (2002), 1337–1372.
- [20] M. E. Taylor, *Pseudodifferential operators*, Princeton University Press, Princeton, N.J., 1981.
- [21] F. Trèves, *Parametrices for a class of Schrödinger equations*, Comm. Pure Appl. Math. **48** (1995), no. 1, 13–78.
- [22] J. Wunsch, *Propagation of singularities and growth for Schrödinger operators*, Duke Math. J. **98** (1999), no. 1, 137–186.
- [23] K. Yajima, *Smoothness and non-smoothness of the fundamental solution of time dependent Schrödinger equations*, Comm. Math. Phys. **181** (1996), no. 3, 605–629.
- [24] S. Zelditch, *Reconstruction of singularities for solutions of Schrödinger’s equation*, Comm. Math. Phys. **90** (1983), no. 1, 1–26.

DEPARTMENT OF MATHEMATICS, AUSTRALIAN NATIONAL UNIVERSITY, CANBERRA, 0200 ACT AUSTRALIA

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON IL USA