

## 547 - Spring 2018 - HW2

January 26, 2018

1. Let  $f, g, h : S^1 \rightarrow X$  be continuous pointed functions. Write down an explicit homotopy between  $(f \cdot g) \cdot h$  and  $f \cdot (g \cdot h)$ .

2. Suppose that  $a : I^1 \rightarrow X$  is a path from  $a(0)$  to  $a(1)$ . Prove that “conjugation by  $a$ ”,

$$f \mapsto a^{-1} \cdot f \cdot a,$$

gives a well defined isomorphism  $\pi_1(X, a(0)) \cong \pi_1(X, a(1))$ .

3. Let  $f, g : X \rightarrow Y$  be pointed maps. If  $f$  is homotopic to  $g$ , then  $f_* = g_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$ .

4. Prove that  $\pi_1$  induces a functor  $\text{Ho}(\mathcal{T}_*) \rightarrow \text{Groups}$ .

5. Describe pushouts and pullbacks in the category  $\text{Ab}$  of abelian groups.

6. Describe pushouts and pullbacks in the category  $\text{CAlg}$  of commutative  $\mathbb{Z}$ -algebras (i.e., commuting rings).

7. Does the forgetful functor  $\text{Ab} \leftarrow \text{CAlg}$  preserve pullbacks? What about pushouts?

8. Prove that  $\pi_1(\mathbb{P}^2(\mathbb{C})) = 0$ . You can do this using van Kampen’s theorem.

9. Do Hatcher, Exercise 1.1.16.

10. Do Hatcher, Exercise 1.1.18.